

Decay of Correlations for the Rauzy–Veech–Zorich Induction Map on the Space of Interval Exchange Transformations and the Central Limit Theorem for the Teichmüller Flow on the Moduli Space of Abelian Differentials.

Alexander I. Bufetov*

February 1, 2008

1 Introduction

The aim of this paper is to prove a stretched-exponential bound for the decay of correlations for the Rauzy–Veech–Zorich induction map on the space of interval exchange transformations (Theorem 4). A Corollary is the Central Limit Theorem for the Teichmüller flow (Theorem 10).

The proof of Theorem 4 proceeds by approximating the induction map by a Markov chain satisfying the Doeblin condition, the method of Sinai [13] and Bunimovich–Sinai [14]. The main “loss of memory” estimate is Lemma 4.

1.1 Interval exchange transformations.

Let m be a positive integer. Let π be a permutation on m symbols. The permutation π will always be assumed *irreducible*, which means that $\pi\{1, \dots, k\} = \{1, \dots, k\}$ only if $k = m$.

Let λ be a vector in $\mathbb{R}_+^m$, $\lambda = (\lambda_1, \dots, \lambda_m)$, $\lambda_i > 0$ for all i . Denote

$$|\lambda| = \sum_{i=1}^m \lambda_i.$$

Consider the half-open interval $[0, |\lambda|)$. Consider the points $\beta_i = \sum_{j < i} \lambda_j$, $\beta_i^\pi = \sum_{j < \pi^{-1}i} \lambda_{\pi^{-1}j}$.

Denote $I_i = [\beta_i, \beta_{i+1})$, $I_i^\pi = [\beta_i^\pi, \beta_{\pi^{-1}i+1}^\pi)$. The length of I_i is λ_i , whereas the length of I_i^π is $\lambda_{\pi^{-1}i}$.

*Department of Mathematics, Princeton University

Set

$$T_{(\lambda, \pi)}(x) = x + \beta_{\pi i}^\pi - \beta_i \text{ for } x \in I_i.$$

The map $T_{(\lambda, \pi)}$ is called an interval exchange transformation corresponding to (λ, π) .

The map $T_{(\lambda, \pi)}$ is an order-preserving isometry from I_i onto $I_{\pi(i)}^\pi$.

We say that λ is *irrational* if there are no rational relations between $|\lambda|$, $\lambda_1, \lambda_2, \dots, \lambda_{m-1}$.

Theorem 1 (Oseledets([5]), Keane([9])) *Let π be irreducible and λ irrational. Then for any $x \in [0, \sum_{i=1}^m \lambda_i)$, the set $\{T_{(\lambda, \pi)}^n x, n \geq 0\}$ is dense in $[0, \sum_{i=1}^m \lambda_i)$.*

1.2 Rauzy operations a and b .

Let (λ, π) be an interval exchange. Assume that π is irreducible and λ is irrational.

Following Rauzy [6], consider the induced map of (λ, π) on the interval $[0, |\lambda| - \min(\lambda_m, \lambda_{\pi^{-1}(m)})]$. The induced map is again an interval exchange of m intervals. For $i, j = 1, \dots, m$, denote by E_{ij} an $m \times m$ matrix of which the i, j -th element is equal to 1, all others to 0. Let E be the $m \times m$ -identity matrix.

1.2.1 Case a : $\lambda_{\pi^{-1}m} > \lambda_m$.

Define

$$A(a, \pi) = \sum_{i=1}^{\pi^{-1}(m)} E_{ii} + E_{m, \pi^{-1}m+1} + \sum_{i=\pi^{-1}m+1}^m E_{i, i+1}$$

$$a\pi(j) = \begin{cases} \pi j, & \text{if } j \leq \pi^{-1}m; \\ \pi m, & \text{if } j = \pi^{-1}m + 1; \\ \pi(j-1), & \text{other } j. \end{cases}$$

If $\lambda_{\pi^{-1}m} > \lambda_m$, then the induced interval exchange of $T_{(\lambda, \pi)}$ on the interval $[0, \sum_{i \neq m} \lambda_i)$ is $T_{(\lambda', \pi')}$, where $\lambda' = A(a, \pi)^{-1}\lambda$ and $\pi' = a\pi$.

1.2.2 Case b : $\lambda_m > \lambda_{\pi^{-1}m}$.

Define

$$A(b, \pi) = E + E_{m, \pi^{-1}m}$$

$$b\pi(j) = \begin{cases} \pi j, & \text{if } \pi j \leq \pi m; \\ \pi j + 1, & \text{if } \pi m < \pi j < m; \\ \pi m + 1, & \text{if } \pi j = m. \end{cases}$$

If $\lambda_{\pi^{-1}m} < \lambda_m$, then the induced interval exchange of $T_{(\lambda, \pi)}$ on the interval $[0, \sum_{i \neq \pi^{-1}m} \lambda_i)$ is $T_{(\lambda', \pi')}$, where $\lambda' = A(b, \pi)^{-1}\lambda$ and $\pi' = b\pi$.

Note that operations a and b are invertible on the space of permutations, namely, we have:

$$a^{-1}\pi(j) = \begin{cases} \pi(j), & \text{if } j \leq \pi^{-1}(m); \\ \pi(j+1), & \text{if } \pi^{-1}(m)+1 < j < m; \\ \pi(\pi^{-1}(\pi(m)+1)), & \text{if } j = m. \end{cases}$$

$$b^{-1}\pi(j) = \begin{cases} \pi(j), & \text{if } \pi(j) \leq \pi(m) \\ m, & \text{if } j = \pi^{-1}(\pi(m)+1); \\ \pi(j)-1, & \text{if } \pi(j) > \pi(m)+1. \end{cases}$$

For $(\lambda, \pi) \in \Delta(\mathcal{R})$, denote

$$T_{a^{-1}}(\lambda, \pi) = (A(a^{-1}\pi, a)\lambda, a^{-1}\pi), \quad T_{b^{-1}}(\lambda, \pi) = (A(b^{-1}\pi, b)\lambda, b^{-1}\pi). \quad (1)$$

The interval exchange $T_{a^{-1}}(\lambda, \pi)$ is the preimage of (λ, π) under the operation a , and the interval exchange $T_{b^{-1}}(\lambda, \pi)$ is the preimage of (λ, π) under the operation b .


Normalize (dividing by $|\lambda| = \lambda_1 + \dots + \lambda_m$) and set:

$$t_{a^{-1}}(\lambda, \pi) = (\frac{A(a^{-1}\pi, a)\lambda}{|A(a^{-1}\pi, a)\lambda|}, a^{-1}\pi), \quad t_{b^{-1}}(\lambda, \pi) = (\frac{A(b^{-1}\pi, b)\lambda}{|A(b^{-1}\pi, b)\lambda|}, b^{-1}\pi). \quad (2)$$

1.3 Rauzy class and Rauzy graph.

If π is an irreducible permutation, then its *Rauzy class* is the set of all permutations that can be obtained from π by applying repeatedly the operations a and b ; the Rauzy class of the permutation π is denoted $\mathcal{R}(\pi)$. Rauzy class has a natural structure of an oriented labelled graph: namely, the permutations of the Rauzy class are the vertices of the graph, and if $\pi = a\pi'$ then we draw an edge from π to π' and label it by a , and if $\pi = b\pi'$ then we draw an edge from π to π' and label it by b . This labelled graph will be called the *Rauzy graph* of the permutation π .

For example, the Rauzy graph of the permutation (4321) is



For a permutation π , consider the set $\{a^n\pi, n \geq 0\}$. This set forms a cycle in the Rauzy graph which will be called the *a-cycle* of π . Similarly, the set $\{b^n\pi, n \geq 0\}$ will be called the *b-cycle* of π .

1.4 The Rauzy-Veech-Zorich induction.

Denote

$$\begin{aligned}\Delta_{m-1} &= \{\lambda \in \mathbb{R}_+^m : |\lambda| = 1\}, \\ \Delta_\pi^+ &= \{\lambda \in \Delta_{m-1}, \lambda_{\pi^{-1}m} > \lambda_m\}, \Delta_\pi^- = \{\lambda \in \Delta_{m-1}, \lambda_m > \lambda_{\pi^{-1}m}\}, \\ \Delta(\mathcal{R}) &= \Delta_{m-1} \times \mathcal{R}(\pi).\end{aligned}$$

Define a map

$$\mathcal{T} : \Delta(\mathcal{R}) \rightarrow \Delta(\mathcal{R})$$

by

$$\mathcal{T}(\lambda, \pi) = \begin{cases} \left(\frac{A(\pi, a)^{-1}\lambda}{|A(\pi, a)^{-1}\lambda|}, a\pi\right), & \text{if } \lambda \in \Delta_\pi^+; \\ \left(\frac{A(\pi, b)^{-1}\lambda}{|A(\pi, b)^{-1}\lambda|}, b\pi\right), & \text{if } \lambda \in \Delta_\pi^-.\end{cases}$$

Each $(\lambda, \pi) \in \Delta(\mathcal{R})$ has exactly two preimages under the map $\mathcal{T}$, namely, $t_{a^{-1}}(\lambda, \pi)$ and $t_{b^{-1}}(\lambda, \pi)$ (2).

The set $\Delta(\mathcal{R})$ is a finite union of simplices. Let $\mathbf{m}$ be the Lebesgue measure on $\Delta(\mathcal{R})$ normalized in such a way that $\mathbf{m}(\Delta(\mathcal{R})) = 1$.

Theorem 2 (Veech[1]) *The map $\mathcal{T}$ has an infinite conservative ergodic invariant measure, absolutely continuous with respect to Lebesgue measure on $\Delta(\mathcal{R})$.*

From this result Veech [1] derives that almost all (with respect to $\mathbf{m}$) interval exchange transformations are uniquely ergodic.

Denote

$$\Delta^+ = \cup_{\pi' \in \mathcal{R}(\pi)} \Delta_{\pi'}^+, \Delta^- = \cup_{\pi' \in \mathcal{R}(\pi)} \Delta_{\pi'}^-.$$

Following Zorich [4], we define the function $n(\lambda, \pi)$ in the following way.

$$n(\lambda, \pi) = \begin{cases} \inf\{k > 0 : \mathcal{T}^k(\lambda, \pi) \in \Delta^-\}, & \text{if } \lambda \in \Delta_\pi^+; \\ \inf\{k > 0 : \mathcal{T}^k(\lambda, \pi) \in \Delta^+\}, & \text{if } \lambda \in \Delta_\pi^-.\end{cases}$$

Define

$$\mathcal{G}(\lambda, \pi) = \mathcal{T}^{n(\lambda, \pi)}(\lambda, \pi).$$

The map $\mathcal{G}$ will be referred to as *the Rauzy-Veech-Zorich induction map* [6, 1, 4].

For $(\lambda, \pi) \in \Delta(\mathcal{R})$, denote

$$t_{a^{-n}}(\lambda, \pi) = t_{a^{-1}}^n(\lambda, \pi), t_{b^{-n}}(\lambda, \pi) = t_{b^{-1}}^n(\lambda, \pi), T_{a^{-n}}(\lambda, \pi) = T_{a^{-1}}^n(\lambda, \pi), T_{b^{-n}}(\lambda, \pi) = T_{b^{-1}}^n(\lambda, \pi).$$

Under the map $\mathcal{G}$, each interval exchange (λ, π) has countably many preimages:

$$\mathcal{G}^{-1}(\lambda, \pi) = \begin{cases} \{t_{a^{-n}}(\lambda, \pi), n \in \mathbb{N}\}, & \text{if } (\lambda, \pi) \in \Delta^+; \\ \{t_{b^{-n}}(\lambda, \pi), n \in \mathbb{N}\}, & \text{if } (\lambda, \pi) \in \Delta^-. \end{cases}$$

Theorem 3 (Zorich[4]) *The map $\mathcal{G}$ has an ergodic invariant probability measure, absolutely continuous with respect to Lebesgue on $\Delta(\mathcal{R})$.*

Denote this invariant measure by ν ; the probability with respect to ν will be denoted by $\mathbb{P}$.

Let $\rho(\lambda, \pi)$ be the density of ν with respect to the Lebesgue measure $\mathbf{m}$. Zorich [4] showed that for any $\pi \in \mathcal{R}$ there exist two positive rational homogeneous of degree $-m$ functions ρ_π^+ , ρ_π^- such that

$$\rho(\lambda, \pi) = \begin{cases} \rho_\pi^+(\lambda), & \text{if } \lambda \in \Delta_\pi^+; \\ \rho_\pi^-(\lambda), & \text{if } \lambda \in \Delta_\pi^-. \end{cases} \quad (3)$$

Remark. In particular, the invariant density is bounded from below: there exists a positive constant $C(\mathcal{R})$, depending on the Rauzy class only and such that $\rho(\lambda, \pi) \geq C(\mathcal{R})$ for any $(\lambda, \pi) \in \Delta(\mathcal{R})$.

The map $\mathcal{G}$ is not mixing: indeed, from the definition of $\mathcal{G}$, we have

$$\mathcal{G}(\Delta^+) = \Delta^-, \quad \mathcal{G}(\Delta^-) = \Delta^+.$$

Let $\mathcal{B}$ be the Borel σ -algebra on $\Delta(\mathcal{R})$, and let $\mathcal{B}_n = \mathcal{G}^{-n}\mathcal{B}$. We have $\mathcal{B}_{n+2} \subset \mathcal{B}_n$. Recall [23] that *exactness* of the map $\mathcal{G}^2$ means, by definition, that the σ -algebra $\cap_{n=1}^\infty \mathcal{B}_{2n}$ is trivial [23] (in other words, that Kolmogorov's 0-1 law holds for the map $\mathcal{G}^2$.)

Proposition 1 *The map $\mathcal{G}^2 : \Delta^+ \rightarrow \Delta^+$ is exact with respect to $\nu|_{\Delta^+}$.*

This Proposition is proven in Section 4; it implies strong mixing for the map $\mathcal{G}^2$.

1.5 The main result

Introduce a metric on Δ_{m-1} by setting

$$d(\lambda, \lambda') = \log \frac{\max_i \frac{\lambda_i}{\lambda'_i}}{\min_i \frac{\lambda_i}{\lambda'_i}}. \quad (4)$$

Now introduce a metric on $\Delta(\mathcal{R})$ by setting

$$d((\lambda, \pi), (\lambda', \pi')) = \begin{cases} 2 + d(\lambda, \lambda'), & \text{if } \pi \neq \pi'; \\ d(\lambda, \lambda'), & \text{if } \pi = \pi'. \end{cases}$$

For $\alpha > 0$, let H_α be the space of functions $\phi : \Delta(\mathcal{R}) \rightarrow \mathbb{R}$ such that if $d((\lambda, \pi), (\lambda', \pi')) \leq 1$, then $|\phi(\lambda, \pi) - \phi(\lambda', \pi')| \leq C d((\lambda, \pi), (\lambda', \pi'))^\alpha$ for some constant C .

Define

$$C_{H_\alpha}(\phi) = \max_{d((\lambda, \pi), (\lambda', \pi')) \leq 1} \frac{|\phi(\lambda, \pi) - \phi(\lambda', \pi')|}{d((\lambda, \pi), (\lambda', \pi'))^\alpha},$$

The main result of this paper is

Theorem 4 *Let $\mathcal{G} : \Delta(\mathcal{R}) \rightarrow \Delta(\mathcal{R})$ be the Rauzy-Veech-Zorich induction map and let ν be the absolutely continuous invariant measure.*

Let $p > 2$. Then, for any $\alpha > 0$, there exist positive constants C, δ such that for any $\phi \in H_\alpha \cap L_p(\Delta^+(\mathcal{R}), \nu)$ and $\psi \in L_2(\Delta^+(\mathcal{R}), \nu)$ we have

$$|\int \phi \times \psi \circ \mathcal{G}^{2n} d\nu - \int \phi d\nu \int \psi d\nu| \leq C \exp(-\delta n^{1/6}) (C_{H_\alpha}(\phi) + |\phi|_{L_p})(|\psi|_{L_2}).$$

Denote by $\mathcal{N}(0, \sigma)$ the Gaussian distribution with mean 0 and variance σ . By [7, 8, 17], we have

Corollary 1 *Let $\phi \in H_\alpha \cap L_p(\Delta(\mathcal{R})^+, \nu)$, $\int \phi d\nu = 0$. Assume that there does not exist $\psi \in L_2(\Delta(\mathcal{R})^+, \nu)$ such that $\phi = \psi \circ \mathcal{G}^2 - \psi$. Then there exists $\sigma > 0$ such that*

$$\frac{1}{\sqrt{N}} \sum_{n=0}^{N-1} \phi \circ \mathcal{G}^{2n} \xrightarrow{d} \mathcal{N}(0, \sigma) \text{ as } N \rightarrow \infty.$$

1.6 Veech's space of zippered rectangles

A zippered rectangle associated to the Rauzy class $\mathcal{R}$ is a quadruple (λ, h, a, π) , where $\lambda \in \mathbb{R}_+^m, h \in \mathbb{R}_+^m, a \in \mathbb{R}^m, \pi \in \mathcal{R}$, and the vectors h and a satisfy the following equations and inequalities (one introduces auxiliary components $a_0 = h_0 = a_{m+1} = h_{m+1} = 0$, and sets $\pi(0) = 0, \pi^{-1}(m+1) = m+1$):

$$h_i - a_i = h_{\pi^{-1}(\pi(i)+1)} - a_{\pi^{-1}(\pi(i)+1)-1}, i = 0, \dots, m$$

$$h_i \geq 0, i = 1, \dots, m, a_i \geq 0, i = 1, \dots, m-1,$$

$$a_i \leq \min(h_i, h_{i+1}) \text{ for } i \neq m, i \neq \pi^{-1}m,$$

$$a_m \leq h_m, a_m \geq -h_{\pi^{-1}m}, a_{\pi^{-1}m} \leq h_{\pi^{-1}m+1}$$

The area of a zippered rectangle is given by the expression $\lambda_1 h_1 + \dots + \lambda_m h_m$. Following Veech, we denote by $\Omega(\mathcal{R})$ the space of all zippered rectangles, corresponding to a given Rauzy class $\mathcal{R}$ and satisfying the condition

$$\lambda_1 h_1 + \dots + \lambda_m h_m = 1.$$

We shall denote by x an individual zippered rectangle.

Veech further defines a map $\mathcal{U}$ and a flow P^t on the space of zippered rectangles in the following way:

$$P^t(\lambda, h, a, \pi) = (e^t \lambda, e^{-t} h, e^{-t} a, \pi).$$

$$\mathcal{U}(\lambda, h, a, \pi) = \begin{cases} (A^{-1}(a, \pi)\lambda, A^t(a, \pi)h, a', a\pi), & \text{if } (\lambda, \pi) \in \Delta^- \\ (A^{-1}(b, \pi)\lambda, A^t(b, \pi)h, a'', b\pi), & \text{if } (\lambda, \pi) \in \Delta^+, \end{cases}$$

where

$$a'_i = \begin{cases} a_i, & \text{if } j < \pi^{-1}m, \\ h_{\pi^{-1}m} + a_{m-1}, & \text{if } i = \pi^{-1}m, \\ a_{i-1}, & \text{other } i. \end{cases}$$

$$a''_i = \begin{cases} a_i, & \text{if } j < m, \\ -h_{\pi^{-1}m} + a_{\pi^{-1}m-1}, & \text{if } i = m. \end{cases}$$

The map $\mathcal{U}$ is invertible; $\mathcal{U}$ and P^t commute ([1]).
Denote

$$\tau(\lambda, \pi) = (\log(|\lambda| - \min(\lambda_m, \lambda_{\pi^{-1}m})),$$

and for $x \in \Omega(\mathcal{R})$, $x = (\lambda, h, a, \pi)$, write

$$\tau(x) = \tau(\lambda, \pi).$$

Now define

$$\mathcal{Y}(\mathcal{R}) = \{x \in \Omega(\mathcal{R}) : |\lambda| = 1\}.$$

and

$$\Omega_0(\mathcal{R}) = \bigcup_{x \in \mathcal{Y}(\mathcal{R}), 0 \leq t \leq \tau(x)} P^t x.$$

$\Omega_0(\mathcal{R})$ is a fundamental domain for $\mathcal{U}$ and, identifying the points x and $\mathcal{U}x$ in $\Omega_0(\mathcal{R})$, we obtain a natural flow, also denoted by P^t , on $\Omega_0(\mathcal{R})$.

The space $\Omega(\mathcal{R})$ has a natural Lebesgue measure class and so does the transversal $\mathcal{Y}(\mathcal{R})$. Veech [1] has proved the following Theorem.

Theorem 5 *There exists a measure $\mu_{\mathcal{R}}$ on $\Omega(\mathcal{R})$, absolutely continuous with respect to Lebesgue, preserved by both the map $\mathcal{U}$ and the flow P^t and such that $\mu_{\mathcal{R}}(\Omega_0(\mathcal{R})) < \infty$.*

For $x \in \mathcal{Y}(\mathcal{R})$, define

$$\mathcal{S}(x) = \mathcal{U}P^{\tau(x)}(x).$$

The map $\mathcal{S}$ is a lift of $\mathcal{T}$ to the space of zippered rectangles: indeed, if

$$\mathcal{S}(\lambda, h, a, \pi) = (\lambda', h', a', \pi'),$$

then $(\lambda', \pi') = \mathcal{T}(\lambda', \pi')$.

Since $\mathcal{Y}(\mathcal{R})$ is a transversal to the flow, the measure $\mu_{\mathcal{R}}$ induces an absolutely continuous measure $\mu_{\mathcal{R}}^{(1)}$ on $\mathcal{Y}(\mathcal{R})$; since $\mu_{\mathcal{R}}$ is both $\mathcal{U}$ and P^t -invariant, the measure $\mu_{\mathcal{R}}^{(1)}$ is $\mathcal{S}$ -invariant. Since $\mu_{\mathcal{R}}(\Omega_0(\mathcal{R})) < \infty$, the measure $\mu_{\mathcal{R}}^{(1)}$ is conservative; it is, however, infinite (Veech [1]).

Zorich [4] constructed a different section for the flow P^t , for which the restricted measure has finite total mass.

Following Zorich [4], define

$$\Omega^+(\mathcal{R}) = \{x = (\lambda, h, a, \pi) : (\lambda, \pi) \in \Delta^+, a_m \geq 0\}.$$

$$\Omega^-(\mathcal{R}) = \{x = (\lambda, h, a, \pi) : (\lambda, \pi) \in \Delta^-, a_m \leq 0\},$$

$$\mathcal{Y}^+(\mathcal{R}) = \mathcal{Y}(\mathcal{R}) \cap \Omega^+(\mathcal{R}), \quad \mathcal{Y}^-(\mathcal{R}) = \mathcal{Y}(\mathcal{R}) \cap \Omega^-(\mathcal{R}), \quad \mathcal{Y}^\pm(\mathcal{R}) = \mathcal{Y}^+(\mathcal{R}) \cup \mathcal{Y}^-(\mathcal{R}).$$

Take $x \in \mathcal{Y}^\pm(\mathcal{R})$, $x = (\lambda, h, a, \pi)$, and define

$$\mathcal{F}(x) = \mathcal{S}^{n(\lambda, \pi)} x.$$

The map $\mathcal{F}$ is a lift of the map $\mathcal{G}$ to the space of zippered rectangles: if

$$\mathcal{F}(\lambda, h, a, \pi) = (\lambda', h', a', \pi'),$$

then $(\lambda', \pi') = \mathcal{G}(\lambda', \pi')$.

We shall see, moreover, that the map $\mathcal{F}$ can be almost surely (with respect to Lebesgue) identified with the natural extension of the map $\mathcal{G}$ (Section 3).

If $x \in \mathcal{Y}^+$, then $\mathcal{F}(x) \in \mathcal{Y}^-$, and if $x \in \mathcal{Y}^-$, then $\mathcal{F}(x) \in \mathcal{Y}^+$. The map $\mathcal{F}$ is the induced map of $\mathcal{S}$ to the subset $\mathcal{Y}^\pm(\mathcal{R})$.

Since $\mathcal{Y}^\pm(\mathcal{R})$ is a transversal to the flow P^t , the measure $\mu_{\mathcal{R}}$ naturally induces an absolutely continuous measure $\bar{\nu}$ on $\mathcal{Y}^\pm(\mathcal{R})$; since $\mu_{\mathcal{R}}$ is both $\mathcal{U}$ and P^t -invariant, the measure $\bar{\nu}$ is $\mathcal{F}$ -invariant.

Zorich [4] proved

Theorem 6 *The measure $\bar{\nu}$ is finite and ergodic for $\mathcal{F}$.*

Since the map $\mathcal{G}$ is exact (as is shown in Section 4), the map $\mathcal{F}$ satisfies the K -property of Kolmogorov, and, in particular, is strongly mixing. Decay of correlations is proven for the map $\mathcal{F}$ as well.

Introduce a metric on the space of zippered rectangles in the following way. Take two zippered rectangles $x = (\lambda, h, a, \pi)$ and $x' = (\lambda', h', a', \pi')$. Write

$$d((\lambda, h, a), (\lambda', h', a')) = \log \frac{\max_i \frac{\lambda_i}{\lambda'_i}, \frac{h_i}{h'_i}, \frac{|a_i|}{|a'_i|}, \frac{|h_i - a_i|}{|h'_i - a'_i|}}{\min_i \frac{\lambda_i}{\lambda'_i}, \frac{h_i}{h'_i}, \frac{|a_i|}{|a'_i|}, \frac{|h_i - a_i|}{|h'_i - a'_i|}}.$$

Define the metric on $\Omega(\mathcal{R})$ by

$$d(x, x') = \begin{cases} d((\lambda, h, a), (\lambda', h', a')) & \text{if } \pi = \pi' \text{ and } \frac{a_m}{a'_m} > 0; \\ 2 + d((\lambda, h, a), (\lambda', h', a')), & \text{otherwise.} \end{cases}$$

As above, for $\alpha > 0$, let H_α be the space of functions $\phi : \mathcal{Y}^\pm(\mathcal{R}) \rightarrow \mathbb{R}$ such that if $d(x, x') \leq 1$, then $|\phi(x) - \phi(x')| \leq C d(x, x')^\alpha$ for some constant C .

Note that the distance $d(x, x')$ is not defined if $a_i = 0$ or $a'_i = 0$ for some $i = 1, \dots, m$; nothing, therefore, is said about the values of a function from H_α at such points. This does not represent a problem, however, since we only need the space H_α for the Central Limit Theorem, and for for such a result we may deal with functions defined almost everywhere.

Define

$$C_{H_\alpha}(\phi) = \max_{d(x, x') \leq 1} \frac{|\phi(x) - \phi(x')|}{d(x, x')^\alpha}.$$

Theorem 7 *Let $\mathcal{F} : \mathcal{Y}^\pm(\mathcal{R}) \rightarrow \mathcal{Y}^\pm(\mathcal{R})$ be the Rauzy-Veech-Zorich induction map on the space of zippered rectangles and let $\overline{\nu}_{\mathcal{R}}$ be the absolutely continuous invariant probability measure. Let $p > 2$. Then, for any $\alpha > 0$, there exist positive constants C, δ such that for any $\phi, \psi \in H_\alpha \cap L_p(\mathcal{Y}(\mathcal{R}), \overline{\nu}_{\mathcal{R}})$ we have*

$$\left| \int \phi \times \psi \circ \mathcal{F}^{2n} d\overline{\nu}_{\mathcal{R}} - \int \phi d\overline{\nu}_{\mathcal{R}} \int \psi d\overline{\nu}_{\mathcal{R}} \right| \leq C \exp(-\delta n^{1/6}) (C_{H_\alpha}(\phi) + |\phi|_{L_p})(C_{H_\alpha}(\psi) + |\psi|_{L_p})$$

Theorem 7 will be established simultaneously with the Theorem 4. Indeed, the map $\mathcal{F}$ can be almost surely identified with the natural extension of the map $\mathcal{G}$, and the method of Markov approximations of Sinai [13] and Bunimovich-Sinai [14] allows to obtain the decay of correlations for the invertible case simultaneously with that for the noninvertible one.

Since the flow P^t is a special flow over the map $\mathcal{F}$, by the Theorem of Melbourne and Török [15], the decay of correlations for the map $\mathcal{F}$ allows to obtain the Central Limit Theorem for the flow P^t .

Denote by X_t the derivative with respect to the flow P^t .

Theorem 8 *Let $p > 2$ and let $\phi \in H_\alpha(\Omega_0(\mathcal{R})) \cap L_p(\Omega_0(\mathcal{R}), \mu_{\mathcal{R}})$ satisfy $\int \phi d\nu = 0$. Assume that there does not exist $\psi \in L_2(\Omega_0(\mathcal{R}), \mu_{\mathcal{R}})$ such that $\phi = X_t \psi$. Then there exists $\sigma > 0$ such that*

$$\frac{1}{\sqrt{T}} \int_0^T \phi \circ P^t \stackrel{d}{\rightarrow} \mathcal{N}(0, \sigma) \text{ as } T \rightarrow \infty.$$

This Theorem will be proved in Section 16.

1.7 Zippered rectangles and the moduli space of holomorphic differentials.

Let $g \geq 2$ be an integer. Take an arbitrary integer vector $\kappa = (k_1, \dots, k_\sigma)$ such that $k_i > 0$, $k_1 + \dots + k_\sigma = 2g - 2$.

Denote by $\mathcal{M}_\kappa$ the moduli space of Riemann surfaces of genus g endowed with a holomorphic differential of area 1 with singularities of orders $k_1, \dots, k_\sigma$. (the *stratum* in the moduli space of holomorphic differentials). Denote by g_t

the Teichmüller flow on $\mathcal{M}_\kappa$ (see [10], [21], [28], [29]). The flow g_t preserves a natural absolutely continuous probability measure on $\mathcal{M}_\kappa$ ([21], [1], [29]). We denote that measure by μ_κ .

A zippered rectangle naturally defines a Riemann surface endowed with a holomorphic differential of area 1. The orders of the singularities of ω are uniquely defined by the Rauzy class of the permutation π ([1]).

For any $\mathcal{R}$ we thus have a map

$$\pi_{\mathcal{R}} : \Omega_{\mathcal{R}} \rightarrow \mathcal{M}_\kappa,$$

where κ is uniquely defined by $\mathcal{R}$.

Veech [1] proved

Theorem 9 (Veech) *1. The set $\pi_0(\Omega_0(\mathcal{R}))$ is a connected component of $\mathcal{M}_\kappa$. Any connected component of any $\mathcal{M}_\kappa$ has the form $\pi_0(\Omega_0(\mathcal{R}))$ for some $\mathcal{R}$.*

- 2. The map π_0 is finite-to-one and almost everywhere locally bijective.*
- 3. $\pi_0(\mathcal{U}x) = \pi_0(x)$.*
- 4. The flow P^t on $\Omega_0(\mathcal{R})$ projects under π_0 to the Teichmüller flow g_t on the corresponding connected component of $\mathcal{M}_\kappa$.*
- 5. $(\pi_{\mathcal{R}})_*\mu_\kappa = \mu_{\mathcal{R}}$.*

A detailed treatment of the relationship between Rauzy classes, zippered rectangles and connected components is given by M.Kontsevich and A.Zorich in [26].

Say that a function $\psi : \mathcal{M}_\kappa \rightarrow \mathbb{R}$ is *Hölder in the sense of Veech* if there exists a Hölder function $\phi : \Omega_0(\mathcal{R}) \rightarrow \mathbb{R}$ such that $\psi \circ \pi_0 = \phi$.

Remark. This definition has a natural interpretation in terms of cohomological coordinates of Hubbard and Masur [28]. Indeed, under the map π_0 the Veech coordinates on the space of zippered rectangles correspond, upto a linear change of variables, to the cohomological coordinates of Hubbard and Masur. Locally, one can associate a Hilbert metric to those coordinates. A function Hölder in the sense of Veech if and only if it is Hölder with respect to that metric. Note that the thus defined local Hilbert distance between two elements in $\mathcal{M}_\kappa$ majorates the Teichmüller distance between their underlying surfaces. Therefore, if a function $\phi : \mathcal{M}_\kappa \rightarrow \mathbb{R}$ is a lift of a smooth function from the underlying moduli space $\mathcal{M}_g$ of compact surfaces of genus g , then ϕ is Hölder in the sense of Veech.

Denote by $\mathcal{X}_t$ the derivative in the direction of the flow g_t .

Theorem 8 and Theorem 9 imply the following

Theorem 10 *Let $\mathcal{H}$ be a connected component of $\mathcal{M}_\kappa$. Let $p > 2$, and let $\psi \in L_p(\mathcal{H}, \mu_\kappa)$ be Hölder in the sense of Veech and satisfy $\int \phi d\mu_\kappa = 0$. Assume that there does not exist $\psi \in L_2(\mathcal{H}, \mu_\kappa)$ such that $\phi = \mathcal{X}_t\psi$. Then there exists $\sigma > 0$ such that*

$$\frac{1}{\sqrt{T}} \int_0^T \phi \circ g_t dt \xrightarrow{d} \mathcal{N}(0, \sigma) \text{ as } T \rightarrow \infty.$$

1.8 Outline of the Proof of Theorem 4.

First, one takes a subset of the space $\Delta(\mathcal{R})$ such that the induced map of $\mathcal{G}$ is uniformly expanding (namely, the set of all interval exchanges such that the renormalization matrix for them is a fixed matrix all whose elements are positive, see Proposition 4; note that the return map on such a subset is an essential element in Veech's proof of unique ergodicity [1]). Then one estimates the statistics of return times in this subset, in the spirit of Lai-Sang Young [11]. After that, the method of Markov approximations, due to Sinai [13], Bunimovich and Sinai [14], is used to complete the proof.

The paper is organized as follows. In Section 2, we state auxiliary propositions about unimodular matrices. In Section 3, following Veech [1] and Zorich [4], we construct symbolic dynamics for the Rauzy-Veech-Zorich induction map $\mathcal{G}$, compute its transition probabilities in the sense of Sinai [13], and identify the natural extension of $\mathcal{G}$ with $\mathcal{F}$. In Section 4, we establish the exactness of $\mathcal{G}^2$. In Section 6, we state the main Lemma 4, whose proof takes Sections 6 – 10. In the remainder of the paper we apply the Markov approximation method of Sinai [13], Bunimovich and Sinai [14], in order to obtain the decay of correlations for $\mathcal{G}$ and $\mathcal{F}$. In the final Section, we apply the Theorem of Melbourne and Török to obtain the Central Limit Theorem for the Teichmüller flow.

2 Matrices

Let A be an $m \times m$ -matrix with positive entries.

Denote

$$\begin{aligned} |A| &= \sum_{i,j=1}^m A_{ij} \\ col(A) &= \max_{i,j,k} \frac{A_{ij}}{A_{kj}}, \\ row(A) &= \max_{i,j,k} \frac{A_{ij}}{A_{ik}} \end{aligned}$$

Proposition 2 *Let Q be a matrix with positive entries, A a matrix with non-negative entries without zero columns or rows.*

Then all entries of the matrices AQ and QA are positive, and, moreover, we have

$$row(AQ) \leq row(Q), col(QA) \leq col(Q)$$

Corollary 2 *Let Q be a matrix with positive entries, A a matrix with nonnegative entries without zero columns or rows.*

$$\text{row}(QAQ) \leq \text{row}(Q), \text{col}(QAQ) \leq \text{col}(Q)$$

Let A be an $m \times m$ matrix with nonnegative entries and determinant 1. Consider the map $J_A : \Delta_{m-1} \rightarrow \Delta_{m-1}$ given by

$$J_A(\lambda) = \frac{A\lambda}{|A\lambda|}.$$

Then

$$\det DJ_A(\lambda) = \frac{1}{|A\lambda|^m}. \quad (5)$$

Suppose all entries of A are positive; then, for any $\lambda, \lambda' \in \Delta_{m-1}$, we have

$$\text{row}(A)^{-m} \leq \frac{\det DJ_A(\lambda)}{\det DJ_A(\lambda')} \leq \text{row}(A)^m, \quad (6)$$

whence we have the following

Proposition 3 *Let $C \subset \Delta_{m-1}$ and let A be a matrix with positive entries and determinant 1. Then*

$$\text{row}(A)^{-m} \frac{\mathbf{m}(C_1)}{\mathbf{m}(C_2)} \leq \frac{\mathbf{m}(J_A(C_1))}{\mathbf{m}(J_A(C_2))} \leq \text{row}(A)^m \frac{\mathbf{m}(C_1)}{\mathbf{m}(C_2)}.$$

We also note the following well-known Lemma (see, for example, [17]):

Lemma 1 *Suppose all entries of the matrix A are positive. Then the map J_A is uniformly contracting with respect to the Hilbert metric.*

3 Symbolic dynamics for $\mathcal{G}$.

First, following Veech [1] and Zorich [4], we describe a Markov partition and a symbolic dynamics for the map $\mathcal{G}^2$, then we identify almost surely the induction map $\mathcal{F}$ on the space of zippered rectangles with the natural extension of $\mathcal{G}$, and, finally, we compute for $\mathcal{G}$ its transition probabilities in the sense of Sinai [25].

3.1 The alphabet

Let $\pi \in \mathcal{R}$, and let n be a positive integer.

Set

$$\Lambda(a, n, \pi) = \{\lambda : \text{there exists } (\lambda', \pi') \text{ such that } \lambda' \in \Delta_{\pi'}^+ \text{ and } (\lambda, \pi) = t_{a-n}(\lambda', \pi')\}$$

$$\Delta(a, n, \pi) = \{(\lambda, \pi), \lambda \in \Lambda(a, n, \pi)\}$$

In other words, $\Delta(a, n, \pi)$ is the set of interval exchange transformations such that the application of the Zorich induction results in the application of the a -operation n times.

The sets $\Delta(a, n, \pi)$ and $\Delta(a, n', \pi')$ are disjoint unless $n = n'$, $\pi = \pi'$, and

$$\Delta_{\pi}^{-} = \cup_{n=1}^{\infty} \Delta(a, n, \pi)$$

up to a set of measure zero (namely, a union of countably many hyperplanes on which Zorich induction is not defined).

If $\pi' = a^n \pi$, then we have

$$\mathcal{G}\Delta(a, n, \pi) = \Delta_{\pi'}^{+}.$$

Similarly, for $\pi \in \mathcal{R}$, and n a positive integer, set

$$\Lambda(b, n, \pi) = \{\lambda : \text{there exists } (\lambda', \pi') \text{ such that } \lambda' \in \Delta_{\pi'}^{-} \text{ and } (\lambda, \pi) = t_{b^{-n}}(\lambda', \pi')\}.$$

$$\Delta(b, n, \pi) = \{(\lambda, \pi), \lambda \in \Lambda(b, n, \pi)\}.$$

In other words, $\Delta(b, n, \pi)$ is the set of interval exchange transformations such that the application of the Zorich induction results in the application of the b -operation n times.

The sets $\Delta(b, n, \pi)$ and $\Delta(b, n', \pi')$ are disjoint unless $n = n'$, $\pi = \pi'$, and

$$\Delta_{\pi}^{+} = \cup_{n=1}^{\infty} \Delta(b, n, \pi)$$

up to a set of measure zero (namely, a union of countably many hyperplanes on which the Zorich induction is not defined).

If $\pi' = b^n \pi$, then, clearly,

$$\mathcal{G}(\Delta(b, n, \pi)) = \Delta_{\pi'}^{-}.$$

Note that the sets $\Delta(a, n, \pi)$ and $\Delta(b, n', \pi')$ are always disjoint, since we have $\Delta(a, n, \pi) \subset \Delta_{\pi}^{-}$, $\Delta(b, n', \pi') \subset \Delta_{\pi'}^{+}$.

The sets $\Delta(a, n, \pi)$, $\Delta(b, n, \pi)$, for all $n > 0$ and all $\pi \in \mathcal{R}$, form a Markov partition for $\mathcal{G}$.

3.2 Words

Consider the alphabet

$$\mathcal{A} = \{(c, n, \pi), c = a \text{ or } b\}$$

For $w_1 \in \mathcal{A}$, $w_1 = (c_1, n_1, \pi_1)$, we write $c_1 = c(w_1)$, $\pi_1 = \pi(w_1)$, $n_1 = n(w_1)$.

For $w_1, w_2 \in \mathcal{A}$, $w_1 = (c_1, n_1, \pi_1)$, $w_2 = (c_2, n_2, \pi_2)$, define the function $B(w_1, w_2)$ in the following way: $B(w_1, w_2) = 1$ if $c_1^{n_1} \pi_1 = \pi_2$ and $c_1 \neq c_2$ and $B(w_1, w_2) = 0$ otherwise.

Let

$$W_{\mathcal{A},B} = \{w = w_1 \dots w_n, w_i \in \mathcal{A}, B(w_i, w_{i+1}) = 1 \text{ for all } i = 1, \dots, n\}.$$

For $w_1 \in \mathcal{A}$, $w_1 = (c_1, n_1, \pi_1)$, set

$$A(w) = A(c_1, c_1^{-n_1} \pi_1) \dots A(c_1, c_1^{-1} \pi_1) A(c_1, \pi_1),$$

and for $w \in W_{\mathcal{A},B}$, $w = w_1 \dots w_n$, set

$$A(w) = A(w_1) \dots A(w_n).$$

Also, for $w_1 \in \mathcal{A}$, $\pi \in \mathcal{R}$, set $w_1^{-1} \pi = c_1^{-n_1} \pi$, and for $w \in W_{\mathcal{A},B}$, $w = w_1 \dots w_n$, set

$$w^{-1} \pi = w_1^{-1} \dots w_n^{-1} \pi.$$

For $w \in W_{\mathcal{A},B}$, define a map $t_w : \Delta(\mathcal{R}) \rightarrow \Delta(\mathcal{R})$ by

$$t_w(\lambda, \pi) = \left(\frac{A(w)\lambda}{|A(w)\lambda|}, w^{-1} \pi \right)$$

Consider also the map

$$T_w(\lambda, \pi) = (A(w)\lambda, w^{-1} \pi)$$

For $w_1 \in \mathcal{A}$, $w_1 = (c_1, n_1, \pi_1)$, we write $\Delta(w_1) = \Delta(c_1, n_1 \pi_1)$.

For $w \in W_{\mathcal{A},B}$, $w = w_1 \dots w_n$, denote

$$\Delta(w) = t_w(\Delta(\mathcal{R})).$$

Then, by definition,

$$\Delta(w) = \{(\lambda, \pi) : (\lambda, \pi) \in \Delta(w_1), \mathcal{G}(\lambda, \pi) \in \Delta(w_2), \dots, \mathcal{G}^{n-1}(\lambda, \pi) \in \Delta(w_n)\}.$$

Say that $w_1 \in \mathcal{A}$ is compatible with $(\lambda, \pi) \in \Delta(\mathcal{R})$ if

1. either $\lambda \in \Delta_\pi^+$, $c_1 = a$, and $a^{n_1} \pi_1 = \pi$
2. or $\lambda \in \Delta_\pi^-$, $c_1 = b$, and $b^{n_1} \pi_1 = \pi$.

Say that a word $w \in W_{\mathcal{A},B}$, $w = w_1 \dots w_n$ is compatible with (λ, π) if w_n is compatible with (λ, π) .

We can write

$$\mathcal{G}^{-n}(\lambda, \pi) = \{t_w(\lambda, \pi) : |w| = n \text{ and } w \text{ is compatible with } (\lambda, \pi)\}.$$

Suppose that a word $w \in W_{\mathcal{A},B}$ is compatible with both (λ, π) and (λ', π) . Then

$$d(t_w(\lambda, \pi), t_w(\lambda', \pi)) \leq d((\lambda, \pi), (\lambda', \pi')).$$

If, moreover, all entries of the the matrix $A(w)$ are positive, then, by Lemma 1, there exists $\alpha(w)$, $0 < \alpha(w) < 1$, such that

$$d(t_w(\lambda, \pi), t_w(\lambda', \pi)) \leq \alpha(w)d((\lambda, \pi), (\lambda', \pi')).$$

We therefore have

Proposition 4 *Let $w \in W_{\mathcal{A},B}$ be such that all entries of the matrix $A(w)$ are positive. Then the return map of $\mathcal{G}$ on $\Delta(w)$ is uniformly expanding with respect to the Hilbert metric.*

3.3 Sequences

Now let

$$\Omega_{\mathcal{A},B} = \{\omega = \omega_1 \dots \omega_n \dots, \omega_n \in \mathcal{A}, B(\omega_n, \omega_{n+1}) = 1 \text{ for all } n \in \mathbb{N}\}$$

and

$$\Omega_{\mathcal{A},B}^{\mathbb{Z}} = \{\omega = \dots \omega_{-n} \dots \omega_1 \dots \omega_n \dots, \omega_n \in \mathcal{A}, B(\omega_n, \omega_{n+1}) = 1 \text{ for all } n \in \mathbb{Z}\}$$

Denote by σ the shift on both these spaces.

There is a natural map $\Phi : \Delta \rightarrow \Omega_{\mathcal{A},B}$ given by the formula

$$\Phi(\lambda, \pi) = \omega_1 \dots \omega_n \dots$$

if

$$\mathcal{G}^n(\lambda, \pi) \in \Delta(\omega_n)$$

The measure ν projects under Φ to a σ -invariant measure on $\Omega_{\mathcal{A},B}$; probability with respect to that measure will be denoted by $\mathbb{P}$.

For $w \in W_{\mathcal{A},B}$, $w = w_1 \dots w_n$, let

$$C(w) = \{\omega \in \Omega_{\mathcal{A},B} : \omega_1 = w_1, \dots, \omega_n = w_n\}.$$

We have then

$$\Delta(w) = \Phi^{-1}(C(w)).$$

W. Veech [1] has proved the following

Proposition 5 *The map Φ is ν -almost surely bijective.*

We thus obtain a symbolic dynamics for the map $\mathcal{G}$.

3.4 The natural extension.

Consider the natural extension for the map $\mathcal{G}$.

The phase space is the space of sequences of interval exchanges; it will be convenient to number them by negative integers. We set:

$$\overline{\Delta}(\mathcal{R}) = \{\mathbf{x} = \dots(\lambda(-n), \pi(-n)), \dots, (\lambda(0), \pi(0)) \mid \mathcal{G}(\lambda(-n), \pi(-n)) = (\lambda(1-n), \pi(1-n)), n = 1, \dots\}$$

The map $\mathcal{G}$ and the invariant measure ν are extended to $\overline{\Delta}$ in the natural way. We shall still denote the probability with respect to the extended measure by $\mathbb{P}$.

We extend the map Φ to a map

$$\overline{\Phi} : \overline{\Delta} \rightarrow \Omega_{\mathcal{A}, B}^{\mathbb{Z}},$$

$$\overline{\Phi}(\lambda) = \dots \omega_{-n} \dots \omega_0 \dots \omega_n \dots,$$

if $(\lambda(-n), \pi(-n)) \in \Delta(\omega_{-n})$, and $\mathcal{G}^n(\lambda(0), \pi(0)) \in \Delta(\omega_n)$.

Now take a zippered rectangle $x \in \Omega(\mathcal{R})$, $x = (\lambda, h, a, \pi)$. Set $\mathcal{F}^n(x) = (\lambda(n), h(n), a(n), \pi(n))$.

Consider a map

$$\tilde{\Phi} : \mathcal{Y}(\mathcal{R}) \rightarrow \Omega_{\mathcal{A}, B}^{\mathbb{Z}}, \tag{7}$$

given by

$$(\lambda, h, a, \pi) \rightarrow \dots \omega_{-n} \dots \omega_0 \dots \omega_n \dots,$$

where

$$(\lambda(n), \pi(n)) \in \Delta(\omega_n)$$

for all $n \in \mathbb{Z}$.

Under the natural projection $(\lambda, h, a, \pi) \rightarrow (\lambda, \pi)$, the $\mathcal{F}$ -invariant measure $\overline{\nu}$ on $\mathcal{Y}^{\pm}(\mathcal{R})$ is mapped to the $\mathcal{G}$ -invariant measure ν on $\Delta(\mathcal{R})$, whence the measure $\tilde{\Phi}_* \overline{\nu}$ is exactly the probability measure $\mathbb{P}$ on the space of bi-infinite sequences. To complete the identification of the spaces $(\mathcal{Y}^{\pm}(\mathcal{R}), \overline{\nu})$ and $(\Omega_{\mathcal{A}, B}^{\mathbb{Z}}, \mathbb{P})$, it remains to show that almost surely there is at most one zippered rectangle corresponding to a given symbolic sequence.

Proposition 6 *Let $\mathbf{q} \in \mathcal{W}_{\mathcal{A}, B}$ be such that all entries of the matrix $A(\mathbf{q})$ are positive. Let $\omega \in \Omega_{\mathcal{A}, B}^{\mathbb{Z}}$ be such that the word $\mathbf{q}$ occurs infinitely many times in ω . Then there exists at most one zippered rectangle corresponding to ω .*

Proof. Write

$$\omega = \dots \omega_{-n} \dots \omega_0 \dots \omega_n \dots,$$

and let (λ, h, a, π) be a zippered rectangle corresponding to ω ; we want to show that (λ, h, a, π) is uniquely defined by ω .

First, (λ, π) is uniquely defined by the "future" $\omega_0 \dots \omega_n \dots$ of ω .

Denote $w(n) = \omega_{-n} \dots \omega_0$, $(\lambda(-n), h(-n), a(-n), \pi(-n)) = \mathcal{F}^{-n}(\lambda, h, a, \pi)$.

For any n , the interval exchange $(\lambda(-n), \pi(-n))$ corresponds to the symbolic sequence $\omega_{-n} \dots \omega_0 \dots$, and, again, is uniquely defined by that sequence.

By definition of the map $\mathcal{F}$, we have

$$\lambda(-n) = \frac{A(w(n))\lambda}{|A(w(n))\lambda|}, \quad h(-n) = (A(w(n))^t)^{-1}h \cdot |A(w(n))\lambda|.$$

Projectively, therefore, we have

$$\mathbb{R}_+ h \subset A(w(n))^t \mathbb{R}_+^m.$$

Since the subword $\mathbf{q}$ occurs infinitely many times, the intersection

$$\bigcap_{n=1}^{\infty} A(w(n))^t \mathbb{R}_+^m$$

consists of a single line and the vector h is therefore uniquely determined by the condition $\langle \lambda, h \rangle = 1$.

It remains to determine the vector a .

By definition of the map $\mathcal{F}$, for any n there exists an orthogonal matrix $U(-n)$, uniquely determined by ω , and a vector $v(-n)$, uniquely determined by the the vectors $h(-n), \dots, h(0)$ and ω , such that

$$\frac{U(-n)a(-n) + v(-n)}{|A(w(n))\lambda|} = a. \quad (8)$$

Now let n be a moment such that all $\lambda(-n)_i > \frac{1}{100m}$ (there are infinitely many such moments). Then $|a(-n)_i| < 100m$ for all $i = 1, \dots, m$ and, (8) since $|A(w(n))\lambda| \rightarrow \infty$ as $n \rightarrow \infty$, (8) implies that a is also uniquely determined by ω .

The proof is complete.

3.5 Transition probabilities.

Take a sequence $c_1 \dots c_n \dots \in \Omega_{\mathcal{A}, B}$. Following Sinai [25], consider the *transition probability*

$$\mathbb{P}(\omega_1 = c_1 | \omega_2 = c_2, \dots, \omega_n = c_n, \dots) = \lim_{n \rightarrow \infty} \frac{\mathbb{P}(c_1 c_2 \dots c_n)}{\mathbb{P}(c_2 \dots c_n)}.$$

In this subsection, we give a formula for this probability in terms of $(\lambda, \pi) = \Phi^{-1}(c_2 \dots c_n \dots)$.

Assume $w_1 \in \mathcal{A}$ is compatible with (λ, π) .

Denote

$$\mathbb{P}(w_1 | (\lambda, \pi)) = \mathbb{P}((\lambda(-1), \pi(-1)) = t_{w_1}(\lambda(0), \pi(0)) | (\lambda(0), \pi(0)) = (\lambda, \pi)).$$

If $w_1 \in \mathcal{A}$ is compatible with (λ, π) , from the definition of $\mathcal{G}$ and from (5) we have

$$\mathbb{P}(w_1 | (\lambda, \pi)) = \frac{\rho(t_{w_1}(\lambda, \pi))}{\rho(\lambda, \pi) |A(w_1)\lambda|^m} \quad (9)$$

Since the invariant density is a homogeneous function of degree $-m$, we have

$$\rho(T_{w_1}(\lambda, \pi)) = \frac{\rho(t_{w_1}(\lambda, \pi))}{|A(w_1)\lambda|^m},$$

and we can rewrite (9) as follows:

$$\mathbb{P}(w_1 | (\lambda, \pi)) = \frac{\rho(T_{w_1}(\lambda, \pi))}{\rho(\lambda, \pi)} \quad (10)$$

Let $w = w_1 \dots w_n$ be compatible with (λ, π) .

Denote

$$\mathbb{P}(w | (\lambda, \pi)) = \mathbb{P}((\lambda(-k), \pi(-k)) = t_{w_{n-k+1}}(\lambda(1-k), \pi(1-k)), k = 1, \dots, n | (\lambda(0), \pi(0)) = (\lambda, \pi)).$$

From (9), by induction, we have

$$\mathbb{P}(w | (\lambda, \pi)) = \frac{\rho(t_w(\lambda, \pi))}{\rho(\lambda, \pi) |A(w)\lambda|^m} \quad (11)$$

Since the invariant density is a homogeneous function of degree $-m$, we have

$$\rho(T_w(\lambda, \pi)) = \frac{\rho(t_w(\lambda, \pi))}{|A(w)\lambda|^m},$$

and we can rewrite (11) as follows:

$$\mathbb{P}(w | (\lambda, \pi)) = \frac{\rho(T_w(\lambda, \pi))}{\rho(\lambda, \pi)} \quad (12)$$

Corollary 3 *There exists $C > 0$ such that the following is true. Suppose $w \in W_{\mathcal{A}, B}$ is compatible with (λ, π) . Then*

$$\mathbb{P}(w | (\lambda, \pi)) \geq \frac{C}{\rho(\lambda, \pi) |A(w)\lambda|^m}$$

Proof: recall that the invariant density is a positive homogeneous function of degree $-m$ and therefore is bounded from below: there exists $C > 0$ such that $\rho(\lambda, \pi) > C$ for all $(\lambda, \pi) \in \Delta(\mathcal{R})$. In particular, $\rho(t_w(\lambda, \pi)) > C$. Substituting into (11), we obtain the result.

For $\epsilon : 0 < \epsilon < 1$, let

$$\Delta_\epsilon = \{(\lambda, \pi) \in \Delta(\mathcal{R}), \min |\lambda_i| \geq \epsilon\}.$$

For any $\epsilon > 0$ there exists a constant $C(\epsilon)$ such that for any $(\lambda, \pi) \in \Delta_\epsilon$ we have $\rho(\lambda, \pi) < C(\epsilon)$.

Corollary 4 *For any $\epsilon > 0$ there exists $C(\epsilon) > 0$ such that if $(\lambda, \pi) \in \Delta_\epsilon$, then*

$$\mathbb{P}(w | (\lambda, \pi)) \geq \frac{C(\epsilon)}{|A(w)|^m}.$$

4 Proof of the Exactness

First, one notes that the discrete parameter π does not give rise to any period, and then the proof follows the standard pattern [27, 17]: since almost every point of any measurable subset is a density point, bounded distortion estimates of Proposition 3 imply that if the measure of a tail event is positive, then it must be arbitrarily close to 1.

In more detail, observe that there exists an integer M such that for any $n > M$ and for any $\pi, \pi' \in \mathcal{R}$ there exist $k_1, \dots, k_{2n}$ such that $a^{k_1} b^{k_2} \dots a^{k_{2n-1}} b^{k_{2n}} \pi = \pi'$. This follows from connectedness of the Rauzy graph and the fact that for any $\pi \in \mathcal{R}$ there exist n_1, n_2 such that $a^{n_1} \pi = b^{n_2} \pi = \pi$.

Let α_0 be the partition of Δ^+ into Δ_π^+ , $\pi \in \mathcal{R}$, and let α_n be the partition into the cylinders $\Delta(w)$, where $w \in \mathcal{W}_{A,B}$, $|w| = 2n$.

Lemma 2 *There exists $k > 0$ such that the following is true. Suppose $C \subset \Delta^+$, and there exists $\pi \in \mathcal{R}$ such that $\Delta_\pi^+ \subset C$. Then $\mathcal{G}^{2k} C = \Delta^+(\mathcal{R})$.*

This implies

Lemma 3 *There exists $k > 0$ such that the following holds. For any $\varepsilon > 0$ there is $\delta > 0$ such that for any $C \subset \Delta^+(\mathcal{R})$ satisfying $\mathbf{m}(C \triangle \Delta_\pi^+) < \delta$, we have $\mathbf{m}(\mathcal{G}^{2k} C \triangle \Delta^+) < \varepsilon$.*

Now suppose $C \subset \Delta^+$ is a $\mathcal{G}^2$ -tail event, i.e., for any $n > 0$ there exists B_n such that $C = \mathcal{G}^{-2n} B_n$ and $0 < \nu(C) < 1$. Then $\nu(B_n) = \nu(C)$ and, by Lemma 3, we can assume that there exists $\varepsilon > 0$ such that for any $\pi \in \mathcal{R}$, we have

$$\mathbf{m}((\Delta^+ \setminus C) \cap \Delta_\pi^+) \geq \varepsilon \quad (13)$$

Let $\mathbf{q} = q_1 \dots q_l$ be a word such that the matrix $A(\mathbf{q})$ is positive.

For almost any $(\lambda, \pi) \in C$ we have

$$\lim_{n \rightarrow \infty} \frac{\mathbf{m}(\alpha_n(\lambda, \pi) \cap C)}{\mathbf{m}(\alpha_n(\lambda, \pi))} = 1 \quad (14)$$

Now let n be such that $\mathcal{G}^{2n}(\lambda, \pi) \in \Delta(\mathbf{q})$. Denote $(\lambda', \pi') = \mathcal{G}^{2n}(\lambda, \pi)$. Let A be the corresponding renormalization matrix, that is, $\lambda = J_A \lambda'$. Then

$A = A_1 A(\mathbf{q})$ for some (unimodular nonnegative integer) matrix A_1 . We have $\alpha_n(\lambda, \pi) = J_A(\Delta_{\pi'}^+)$. By Proposition 3, from (13), we deduce that there exists ε' , not depending on n such that

$$\frac{\mathbf{m}(\alpha_n(\lambda, \pi) \cap (\Delta^+ \setminus C))}{\mathbf{m}(\alpha_n(\lambda, \pi))} \geq \varepsilon'.$$

Since, by ergodicity, for almost any (λ, π) we can find infinitely many n such that $\mathcal{G}^{2n}(\lambda, \pi) \in \Delta(\mathbf{q})$, we arrive at a contradiction with (14), which gives the exactness of $\mathcal{G}^2$.

5 The Main Lemma

We shall suppose from now on that the Rauzy class $\mathcal{R}$ is fixed and will often suppress it from notation.

For $\epsilon : 0 < \epsilon < 1$, define, in the same way as above,

$$\Delta_\epsilon = \{(\lambda, \pi) \in \Delta(\mathcal{R}), \min |\lambda_i| \geq \epsilon\}.$$

Lemma 4 *There exist positive constants γ, K, p such that the following is true for any $\epsilon > 0$. Suppose $(\lambda, \pi) \in \Delta_\epsilon$. Then*

$$\mathbb{P}\{\exists n \leq K|\log \epsilon|, (\lambda(-n), \pi(-n)) \in \Delta_\gamma | (\lambda(1), \pi(1)) = (\lambda, \pi)\} \geq p.$$

From Corollary 4, we obtain

Corollary 5 *Let $\mathbf{q} \in W_{\mathcal{A}, B}$, $\mathbf{q} = q_1 \dots q_l$ be such that all entries of the matrix $A(\mathbf{q})$ are positive. Then there exist positive constants $K(\mathbf{q}), p(\mathbf{q})$ such that the following is true for any $\epsilon > 0$. Suppose $(\lambda, \pi) \in \Delta_\epsilon$. Then*

$$\mathbb{P}\{\exists n \leq K(\mathbf{q})|\log \epsilon|, (\lambda(-n), \pi(-n)) \in \Delta(\mathbf{q}) | (\lambda(1), \pi(1)) = (\lambda, \pi)\} \geq p(\mathbf{q}).$$

Informally, the proof of Lemma 4 proceeds by getting rid of small intervals. For $\gamma > 0$, $k \leq m$, denote

$$\Delta_{\gamma, k} = \{(\lambda, \pi) : \exists i_1, \dots, i_k : \lambda_{i_1}, \dots, \lambda_{i_k} \geq \gamma\}.$$

and

$$\Delta_{\gamma, k, \epsilon} = \{(\lambda, \pi) : \lambda_i \geq \epsilon \text{ for all } i = 1, \dots, m \text{ and } \exists i_1, \dots, i_k : \lambda_{i_1}, \dots, \lambda_{i_k} \geq \gamma\}.$$

Lemma 4 follows from

Lemma 5 *There exist constants L, K, p , depending only on the Rauzy class, such that the following is true for any γ, k, ϵ .*

Assume $(\lambda, \pi) \in \Delta_{\gamma, k, \epsilon}$.

Then

$$\mathbb{P}\{\exists n \leq K|\log \epsilon| : (\lambda(-n), \pi(-n)) \in \Delta_{\gamma/L, k+1, \epsilon/L} | (\lambda(1), \pi(1)) = (\lambda, \pi)\} \geq p.$$

Lemma 5 is proved in the next four sections.

6 An estimate on the number of Rauzy operations.

Recall that, if $(\lambda, \pi) \in \Delta^+$, then the $\mathcal{G}$ -preimages of (λ, π) are the exchanges $t_{a^{-n}}(\lambda, \pi)$, $n = 1, \dots$ whereas if $(\lambda, \pi) \in \Delta^-$, then the $\mathcal{G}$ -preimages of (λ, π) are the exchanges $t_{b^{-n}}(\lambda, \pi)$, $n = 1, \dots$

Denote

$$\mathbf{p}_n(\lambda, \pi) = \begin{cases} \mathbb{P}((\lambda(-1), \pi(-1)) = t_{a^{-n}}(\lambda, \pi) | (\lambda(0), \pi(0)) = (\lambda, \pi)), & \text{if } (\lambda, \pi) \in \Delta^+; \\ \mathbb{P}((\lambda(-1), \pi(-1)) = t_{b^{-n}}(\lambda, \pi) | (\lambda(0), \pi(0)) = (\lambda, \pi)), & \text{if } (\lambda, \pi) \in \Delta^-. \end{cases}$$

For $\lambda \in \mathbb{R}_+^m$, set

$$T_{a^{-1}}^{(\pi)}(\lambda) = A(a^{-1}\pi, a)\lambda, \quad t_{a^{-1}}^{(\pi)}(\lambda) = \frac{A(a^{-1}\pi, a)\lambda}{|A(a^{-1}\pi, a)\lambda|},$$

$$T_{b^{-1}}^{(\pi)}(\lambda) = A(b^{-1}\pi, b)\lambda, \quad t_{b^{-1}}^{(\pi)}(\lambda) = \frac{A(b^{-1}\pi, b)\lambda}{|A(b^{-1}\pi, b)\lambda|},$$

and

$$\begin{aligned} T_{a^{-n}}^{(\pi)}(\lambda) &= T_{a^{-1}}^{(a^{1-n}\pi)} \dots T_{a^{-1}}^{(\pi)}\lambda, \quad t_{a^{-n}}^{(\pi)}(\lambda) = t_{a^{-1}}^{(a^{1-n}\pi)} \dots t_{a^{-1}}^{(\pi)}\lambda, \\ T_{b^{-n}}^{(\pi)}(\lambda) &= T_{b^{-1}}^{(b^{1-n}\pi)} \dots T_{b^{-1}}^{(\pi)}\lambda, \quad t_{b^{-n}}^{(\pi)}(\lambda) = t_{b^{-1}}^{(b^{1-n}\pi)} \dots t_{b^{-1}}^{(\pi)}\lambda, \end{aligned}$$

so that we have

$$\begin{aligned} t_{a^{-n}}(\lambda, \pi) &= (t_{a^{-n}}^{(\pi)}\lambda, a^{-n}\pi), \quad T_{a^{-n}}(\lambda, \pi) = (T_{a^{-n}}^{(\pi)}\lambda, a^{-n}\pi), \\ t_{b^{-n}}(\lambda, \pi) &= (t_{b^{-n}}^{(\pi)}\lambda, b^{-n}\pi), \quad T_{b^{-n}}(\lambda, \pi) = (T_{b^{-n}}^{(\pi)}\lambda, b^{-n}\pi). \end{aligned}$$

Lemma 6 *If $(\lambda, \pi) \in \Delta^+$, then, for any $N \geq 1$, we have*

$$\sum_{n=N+1}^{\infty} \mathbf{p}_n(\lambda, \pi) = \frac{\rho_{a^{-N}\pi}^+(T_{a^{-N}}^{(\pi)}(\lambda))}{\rho_{\pi}^+(\lambda)}$$

If $(\lambda, \pi) \in \Delta^-$, then, for any $N \geq 1$, we have

$$\sum_{n=N+1}^{\infty} \mathbf{p}_n(\lambda, \pi) = \frac{\rho_{b^{-N}\pi}^-(T_{b^{-N}}^{(\pi)}(\lambda))}{\rho_{\pi}^-(\lambda)}$$

Proof: We only consider the case $(\lambda, \pi) \in \Delta^+$. In this case, the formula (10) can be written as

$$\mathbf{p}_n(\lambda, \pi) = \frac{\rho_{a^{-n}\pi}^-(T_{a^{-n}}^{(\pi)}\lambda)}{\rho_{\pi}^+(\lambda)},$$

whence we can write

$$\rho_{\pi}^+(\lambda) = \sum_{n=1}^{\infty} \rho_{a^{-n}\pi}^-(T_{a^{-n}}^{(\pi)}\lambda). \quad (15)$$

Note that this formula is true for *any* permutation π and *any* λ (i.e., even if $\lambda \notin \Delta_{\pi}^+$, the formula, being an identity between rational functions, still holds). Since, for any λ , we have

$$T_{a^{-n-N}}^{(\pi)}\lambda = T_{a^{-n}}^{(a^{-N}\pi)}(T_{a^{-N}}^{(\pi)}\lambda),$$

from (15) we obtain

$$\rho_{a^{-N}\pi}^+(T_{a^{-N}}^{(\pi)}\lambda) = \sum_{n=1}^{\infty} \rho_{a^{-n-N}\pi}^-(T_{a^{-n-N}}^{(\pi)}\lambda) = \rho_{\pi}^+(\lambda) \left(\sum_{n=N+1}^{\infty} \mathbf{p}_n(\lambda, \pi) \right),$$

and the Lemma is proved.

6.1 Bounded growth

Let $(\lambda, \pi) \in \Delta(\mathcal{R})$.

Define

$$(\lambda^{(n)}, \pi^{(n)}) = \begin{cases} t_{a^{-n}}(\lambda, \pi), & \text{if } (\lambda, \pi) \in \Delta^+; \\ t_{b^{-n}}(\lambda, \pi), & \text{if } (\lambda, \pi) \in \Delta^-. \end{cases}$$

$$(\Lambda^{(n)}, \pi^{(n)}) = \begin{cases} T_{a^{-n}}(\lambda, \pi), & \text{if } (\lambda, \pi) \in \Delta^+; \\ T_{b^{-n}}(\lambda, \pi), & \text{if } (\lambda, \pi) \in \Delta^-. \end{cases}$$

We have

$$\mathcal{G}^{-1}(\lambda, \pi) = \{(\lambda^{(n)}, \pi^{(n)}), n = 1, \dots\}.$$

and

$$\mathbf{p}_n = \mathbb{P}((\lambda(-1), \pi(-1)) = (\lambda^{(n)}, \pi^{(n)}) | ((\lambda(0), \pi(0)) = (\lambda, \pi)).$$

For any $n \in \mathbb{N}$, there exists $i(n) \in \{1, \dots, m\}$ such that

$$|\Lambda^{(n)}| - |\Lambda^{(n-1)}| = \lambda_{i(n)}.$$

If $(\lambda(-1), \pi(-1))$ is a $\mathcal{G}$ -preimage of (λ, π) and $(\lambda(-1), \pi(-1)) = t_{c^{-n}}(\lambda, \pi)$, $c = a$ or b , then we define a vector $\Lambda(-1)$ by the relation $(\Lambda(-1), \pi(-1)) = T_{c^{-n}}(\lambda, \pi)$ (in other words, $(\Lambda(-1), \pi(-1))$ is the Zorich preimage without normalization).

Lemma 7 *There exists a constant $C(\mathcal{R})$, depending on the Rauzy class only, such that for any $(\lambda, \pi) \in \Delta(\mathcal{R})$ we have*

$$\mathbb{P}(|\Lambda(-1)| > K | (\lambda(0), \pi(0)) = (\lambda, \pi)) < \frac{C(\mathcal{R})}{K-2}.$$

For definiteness, assume $\lambda \in \Delta_{\pi}^{-}$ (the proof is completely identical in the other case). Then $\mathcal{G}$ -preimages of (λ, π) are $(\lambda^{(n)}, \pi^{(n)}) = t_{b^{-n}}(\lambda, \pi)$, $n = 1, 2, \dots$.

By construction [4], the invariant density ρ_{π}^{-} has the form

$$\rho_{\pi}^{-}(\lambda) = \sum_{i=1}^N \frac{1}{l_{i1}(\lambda) l_{i2}(\lambda) \dots l_{im}(\lambda)},$$

where the functions l_{ij} are linear:

$$l_{ij}(\lambda) = a_{ij}^{(1)} \lambda_1 + \dots + a_{ij}^{(m)} \lambda_m,$$

and all $a_{ij}^{(r)}$ are nonnegative (in fact, $a_{ij}^{(r)} = 0$ or 1 , but we do not need this fact here).

Let l be the length of the a -cycle of π , that is, the smallest such number that $a^l \pi = \pi$.

Since for any $k > 0$ we have $a^{-kl} \pi = \pi$, from Lemma 6 we obtain

$$\sum_{n=kl+1}^{\infty} \mathbf{p}_n(\lambda, \pi) = \frac{\rho_{\pi}^{-}(\Lambda^{(kl)})}{\rho_{\pi}^{-}(\lambda)}.$$

As noted above, for any $n > 0$ there exists $\lambda_{i(n)}$ such that

$$|\Lambda^{(n)}| - |\Lambda^{(n-1)}| = \lambda_{i(n)},$$

and, in fact,

$$\Lambda^{(n)} = (\lambda_1, \dots, \lambda_{m-1}, \lambda_m + \lambda_{i(1)} + \dots + \lambda_{i(n)}).$$

Since

$$\sum_{n=kl+1}^{\infty} \mathbf{p}_n(\lambda, \pi) \rightarrow 0 \text{ as } k \rightarrow \infty,$$

for any $i = 1, \dots, N$ there exists j such that $a_{ij}^{(m)} > 0$. Renumbering, if necessary, the linear forms l_{ij} , we may assume that $a_{i1}^{(m)} > 0$ for any i . Denote $\epsilon = \min a_{i1}^{(m)}$ and $L = \max a_{i1}^{(r)}$. For any $\lambda \in \mathbb{R}_+^m$ we have then

$$\epsilon \lambda_m \leq l_{i1}(\lambda) \leq L |\lambda|,$$

whence

$$\frac{\rho_{\pi}^{-}(\Lambda^{(kl)})}{\rho_{\pi}^{-}(\lambda)} \leq \frac{L}{\epsilon(\lambda_m + \lambda_{i(1)} + \dots + \lambda_{i(kl)})}. \quad (16)$$

Let N be the smallest number such that $|\Lambda(-N)| > K$ and let s be the largest such integer that $sl < N$. Then $|\Lambda(-sl)| > K - 1$ (because all $\lambda_{i(sl+1)}, \dots, \lambda_{i(N)}$ are all distinct) and $\lambda_m + \lambda_{i(1)} + \dots + \lambda_{i(sl)} > K - 2$ (because $|\Lambda(-sl)| = 1 + \lambda_{i(1)} + \dots + \lambda_{i(sl)}$).

Therefore, by (16), we obtain

$$\frac{\rho_{\pi}^{-}(\Lambda^{(kl)})}{\rho_{\pi}^{-}(\lambda)} \leq \frac{L}{\epsilon} \frac{1}{K - 2},$$

and the Lemma is proved.

Lemma 8 Suppose $(\lambda, \pi) \in \Delta^+$, and let l be the length of the a -cycle of π .

Then, for any $k \geq 1$, we have

$$\sum_{n=kl+1}^{\infty} \mathbf{p}_n(\lambda, \pi) \geq \left(\frac{\lambda_{\pi^{-1}m}}{\lambda_{\pi^{-1}m} + k} \right)^m.$$

Suppose $(\lambda, \pi) \in \Delta^-$, and let l be the length of the b -cycle of π .

Then, for any $k \geq 1$, we have

$$\sum_{n=kl+1}^{\infty} \mathbf{p}_n(\lambda, \pi) \geq \left(\frac{\lambda_m}{\lambda_m + k} \right)^m.$$

Proof. Again, we only consider the case $(\lambda, \pi) \in \Delta^-$, as the proof of the other case is identical.

$$\sum_{n=kl+1}^{\infty} \mathbf{p}_n(\lambda, \pi) = \frac{\rho_{\pi}^{-}(\Lambda^{(kl)})}{\rho_{\pi}^{-}(\lambda)}.$$

Set $\Lambda^{(kl)} = (\Lambda_1^{(kl)}, \dots, \Lambda_m^{(kl)})$.

For $k = 1$ we have $\Lambda_i^{(l)} = \lambda_i$ for $i < m$ and $\Lambda_m^{(l)} = \lambda_m + \lambda_{i(1)} + \dots + \lambda_{i(l)}$, and for arbitrary k by induction we obtain $\Lambda_i^{(kl)} = \lambda_i$ for $i < m$ and $\Lambda_m^{(kl)} = \lambda_m + k(\lambda_{i(1)} + \dots + \lambda_{i(l)})$.

Note that $\lambda_{i(1)} + \dots + \lambda_{i(l)} \leq 1$ (since $i(1), \dots, i(l)$ are all distinct).

As in the proof of the previous Lemma, write

$$\rho_{\pi}^{-}(\lambda) = \sum_{i=1}^N \frac{1}{l_{i1}(\lambda)l_{i2}(\lambda) \dots l_{im}(\lambda)},$$

whence

$$\frac{\rho_{\pi}^{-}(\Lambda^{(kl)})}{\rho_{\pi}^{-}(\lambda)} \geq \min_i \frac{l_{i1}(\lambda)l_{i2}(\lambda) \dots l_{im}(\lambda)}{l_{i1}(\Lambda^{(kl)})l_{i2}(\Lambda^{(kl)}) \dots l_{im}(\Lambda^{(kl)})}. \quad (17)$$

For any linear form $l(\lambda) = a_1\lambda_1 + \cdots + a_m\lambda_m$, $a_i \geq 0$, we have

$$\frac{l(\Lambda^{(kl)})}{l(\lambda)} \geq \frac{\lambda_m}{\lambda_m + k(\lambda_{i(1)} + \cdots + \lambda_{i(l)})} \geq \frac{\lambda_m}{\lambda_m + k},$$

and the Lemma follows.

7 An estimate on the probability of stopping.

Lemma 9 *For any $\gamma > 0$, there exists $c(\gamma) > 0$ such that if $\lambda_{i(N)} > \gamma$, then*

$$\frac{\mathbf{p}_N(\lambda, \pi)}{\sum_{n=N+1}^{\infty} \mathbf{p}_n(\lambda, \pi)} \geq c(\gamma)$$

From Lemma 8 we immediately have the following Corollary.

Corollary 6 *For any $\gamma > 0$, there exists $c(\gamma) > 0$ such that the following is true.*

Assume $(\lambda, \pi) \in \Delta^+$, $\lambda_{i(N)} > \gamma$, $\lambda_{\pi^{-1}m} > \gamma$. Then

$$\mathbf{p}_N \geq \frac{c(\gamma)}{N^m}.$$

Similarly, if $(\lambda, \pi) \in \Delta^-$, $\lambda_{i(N)} > \gamma$, $\lambda_m > \gamma$, then

$$\mathbf{p}_N \geq \frac{c(\gamma)}{N^m}.$$

If $(\lambda, \pi) \in \Delta^+$, then, by the definition of $\mathbf{p}_n(\lambda, \pi)$ and by Lemma 6, we have

$$\begin{aligned} \mathbf{p}_N(\lambda, \pi) &= \frac{\rho_{a^{-N}\pi}^-(T_{a^{-N}}^{(\pi)}\lambda)}{\rho_{\pi}^+(\lambda)}, \\ \sum_{n=N+1}^{\infty} \mathbf{p}_n(\lambda, \pi) &= \frac{\rho_{a^{-N}\pi}^+(T_{a^{-N}}^{(\pi)}(\lambda))}{\rho_{\pi}^+(\lambda)}, \end{aligned}$$

and, therefore,

$$\frac{\mathbf{p}_N(\lambda, \pi)}{\sum_{n=N+1}^{\infty} \mathbf{p}_n(\lambda, \pi)} = \frac{\rho_{a^{-N}\pi}^-(T_{a^{-N}}^{(\pi)}\lambda)}{\rho_{a^{-N}\pi}^+(T_{a^{-N}}^{(\pi)}(\lambda))}.$$

Lemma 9 follows now from the following

Lemma 10 *For any $\gamma > 0$ there exists a constant $c(\gamma) > 0$ such that the following is true. Let $(\lambda, \pi) \in \Delta(\mathcal{R})$. If $\lambda_{\pi^{-1}m+1} > \gamma$, then*

$$\frac{\rho_{\pi}^-(\lambda)}{\rho_{\pi}^+(\lambda)} \geq c(\gamma).$$

If $\lambda_{\pi^{-1}(\pi(m)+1)} > \gamma$, then

$$\frac{\rho_{\pi}^+(\lambda)}{\rho_{\pi}^-(\lambda)} \geq c(\gamma).$$

The proof of Lemma 10 will take the remainder of this section.

First, we modify Veech's coordinates on the space of zippered rectangles. Take a zippered rectangle $(\lambda, h, a, \pi) \in \Delta(\mathcal{R})$, and introduce the vector $\delta = (\delta_1, \dots, \delta_m) \in \mathbb{R}^m$ by the formula

$$\delta_i = a_{i-1} - a_i, \quad i = 1, \dots, m$$

(here we assume, as always, $a_0 = a_{m+1} = 0$).

Proposition 7 *The data (λ, π, δ) determine the zippered rectangle (λ, h, a, π) uniquely.*

Remark. The coordinates (λ, π, δ) on the space of zippered rectangles have a natural interpretation in terms of the cohomological coordinates of Hubbard and Masur [28]: namely, the λ_i are the real parts of the corresponding cycles, and the δ_i are (minus) the imaginary parts.

Proof of Proposition 7. For any $i = 1, \dots, m$, we have

$$a_i = -\delta_1 - \dots - \delta_i, \quad (18)$$

so the vector a is uniquely defined by δ . It remains to show that the vector h is uniquely defined by δ , and, to do this, we shall express the h through the a . First note that

$$h_{\pi^{-1}m} = a_{\pi^{-1}m} - a_m.$$

Now, if $i \neq \pi^{-1}m$, then $i = \pi^{-1}(k-1)$ for some $k \in \{1, \dots, m\}$. The equation

$$h_i - a_i = h_{\pi^{-1}(\pi(i)+1)} - a_{\pi^{-1}(\pi(i)+1)-1}. \quad (19)$$

then takes the form

$$h_{\pi^{-1}(k-1)} - a_{\pi^{-1}(k-1)} = h_{\pi^{-1}(k)} - a_{\pi^{-1}(k)-1},$$

or, equivalently,

$$h_{\pi^{-1}(k)} = a_{\pi^{-1}(k)-1} + h_{\pi^{-1}(k-1)} - a_{\pi^{-1}(k-1)}.$$

Since

$$h_{\pi^{-1}1} = a_{\pi^{-1}1-1},$$

by induction, we obtain

$$h_{\pi^{-1}k} = a_{\pi^{-1}k-1} + \sum_{l=1}^{k-1} (a_{\pi^{-1}l-1} - a_{\pi^{-1}l})$$

for any $k = 1, \dots, m$, and the Lemma is proved.

The above computations give us the following expression for h in terms of δ :

$$h_{\pi^{-1}k} = - \sum_{i=1}^{\pi^{-1}k-1} \delta_i + \sum_{l=1}^{k-1} \delta_{\pi^{-1}(l)} \quad (20)$$

or, equivalently,

$$h_r = -\sum_{i=1}^{r-1} \delta_i + \sum_{l=1}^{\pi(r)-1} \delta_{\pi^{-1}l}. \quad (21)$$

Rewriting the inequalities defining the zippered rectangle in terms of δ , we obtain by a straightforward computation the following system:

$$\delta_1 + \cdots + \delta_i \leq 0, \quad i = 1, \dots, m-1.$$

$$\delta_{\pi^{-1}1} + \cdots + \delta_{\pi^{-1}i} \geq 0, \quad i = 1, \dots, m-1.$$

The parameter $a_m = -(\delta_1 + \cdots + \delta_m)$ can be both positive and negative. Introduce the following cones in $\mathbb{R}^m$:

$$K_\pi = \{\delta = (\delta_1, \dots, \delta_m) : \delta_1 + \cdots + \delta_i \leq 0, \delta_{\pi^{-1}1} + \cdots + \delta_{\pi^{-1}i} \geq 0, i = 1, \dots, m-1\},$$

$$K_\pi^+ = K_\pi \cap \{\delta : \sum_{i=1}^m \delta_i \leq 0\}, K_\pi^- = K_\pi \cap \{\delta : \sum_{i=1}^m \delta_i \geq 0\}.$$

We have established the following

Proposition 8 *For $(\lambda, \pi) \in \Delta(\mathcal{R})$ and an arbitrary $\delta \in K_\pi$ there exists a unique zippered rectangle (λ, h, a, π) corresponding to the parameters (λ, π, δ) .*

In what follows, we shall simply refer to the zippered rectangle (λ, π, δ) .

Remark. It would be interesting to write down explicitly the generating vectors for the cones K_π , K_π^+ , K_π^- ; in particular, that would allow to give an explicit expression for the invariant densities of Veech [1] and Zorich [4].

Denote by $Area(\lambda, \pi, \delta)$ the area of the zippered rectangle (λ, π, δ) . We have:

$$\begin{aligned} Area(\lambda, \pi, \delta) &= \sum_{r=1}^m \lambda_r h_r = \sum_{r=1}^m \lambda_r \left(-\sum_{i=1}^{r-1} \delta_i + \sum_{l=1}^{\pi(r)-1} \delta_{\pi^{-1}l} \right) = \\ &= \sum_{i=1}^m \delta_i \left(-\sum_{r=i+1}^m \lambda_r + \sum_{r=\pi(i)+1}^m \lambda_{\pi^{-1}r} \right) = 1. \end{aligned} \quad (22)$$

A straightforward computation shows that in the coordinates (λ, π, δ) the Rauzy induction map is written as follows:

$$\mathcal{T}(\lambda, \pi, \delta) = \begin{cases} \left(\left(\frac{A(\pi, b)^{-1} \lambda}{|A(\pi, b)^{-1} \lambda|} \right), b\pi, A(\pi, b)^{-1} \delta \cdot |A(\pi, b)^{-1} \lambda| \right), & \text{if } \lambda \in \Delta_\pi^+; \\ \left(\left(\frac{A(\pi, a)^{-1} \lambda}{|A(\pi, a)^{-1} \lambda|} \right), a\pi, A(\pi, a)^{-1} \delta \cdot |A(\pi, a)^{-1} \lambda| \right), & \text{if } \lambda \in \Delta_\pi^-. \end{cases}$$

For $\lambda \in \mathbb{R}_+^m$, denote

$$\begin{aligned}
K(\lambda, \pi) &= K_\pi \cap \{\delta : \text{Area}(\lambda, \pi, \delta) \leq 1\}, \\
K^+(\lambda, \pi) &= K_\pi^+ \cap \{\delta : \text{Area}(\lambda, \pi, \delta) \leq 1\}, \\
K^-(\lambda, \pi) &= K_\pi^- \cap \{\delta : \text{Area}(\lambda, \pi, \delta) \leq 1\}.
\end{aligned}$$

Denote by vol_m the Lebesgue measure in $\mathbb{R}^m$.
Set

$$\mathbf{r}(\lambda, \pi) = \text{vol}_m(K(\lambda, \pi)), \mathbf{r}^+(\lambda, \pi) = \text{vol}_m(K^+(\lambda, \pi)), \mathbf{r}^-(\lambda, \pi) = \text{vol}_m(K^-(\lambda, \pi)).$$

By definition, the functions $\mathbf{r}, \mathbf{r}^+, \mathbf{r}^-$ are positive rational functions, homogeneous of degree $-m$.

Lemma 11 1. $\mathbf{r}^-(\lambda, \pi) = \mathbf{r}(T_{b^{-1}}(\lambda, \pi))$.

$$2. \mathbf{r}^+(\lambda, \pi) = \mathbf{r}(T_{a^{-1}}(\lambda, \pi)).$$

$$3. \mathbf{r}(\lambda, \pi) = \mathbf{r}(T_{a^{-1}}(\lambda, \pi)) + \mathbf{r}(T_{b^{-1}}(\lambda, \pi)).$$

Proof. If

$$\delta = (\delta_1, \dots, \delta_m) \in K^-(\lambda, \pi),$$

then

$$\tilde{\delta} = (\delta_1, \dots, \delta_{m-1}, \delta_m + \delta_{\pi^{-1}m}) \in K(T_{b^{-1}}(\lambda, \pi)),$$

and vice versa. This gives a volume-preserving bijection between $K^-(\lambda, \pi)$ and $K(T_{b^{-1}}(\lambda, \pi))$, whence $\mathbf{r}^-(\lambda, \pi) = \mathbf{r}(T_{b^{-1}}(\lambda, \pi))$. The second assertion is proved in the same way, and the third follows from the first two.

Corollary 7

$$\begin{aligned}
\mathbf{r}^+(\lambda, \pi) &= \sum_{n=1}^{\infty} \mathbf{r}^-(T_{a^{-n}}(\lambda, \pi)). \\
\mathbf{r}^-(\lambda, \pi) &= \sum_{n=1}^{\infty} \mathbf{r}^+(T_{b^{-n}}(\lambda, \pi)).
\end{aligned}$$

We only prove the first assertion. We have

$$\mathbf{r}^+(\lambda, \pi) = \mathbf{r}(T_{a^{-1}}(\lambda, \pi)) = \mathbf{r}^+(T_{a^{-1}}(\lambda, \pi)) + \mathbf{r}^-(T_{a^{-1}}(\lambda, \pi)) = \mathbf{r}(T_{a^{-2}}(\lambda, \pi)) + \mathbf{r}^-(T_{a^{-1}}(\lambda, \pi)).$$

Proceeding by induction,

$$\mathbf{r}^+(\lambda, \pi) = \sum_{n=1}^N \mathbf{r}^-(T_{a^{-n}}(\lambda, \pi)) + \mathbf{r}(T_{a^{-N-1}}(\lambda, \pi)).$$

Since

$$T_{a^{-N-1}}(\lambda, \pi) = (T_{a^{-N-1}}^{(\pi)}(\lambda), a^{-N-1}\pi),$$

and $|T_{a^{-N-1}}^{(\pi)}(\lambda)| \rightarrow \infty$ as $N \rightarrow \infty$, we obtain $\mathbf{r}(T_{a^{-N-1}}(\lambda, \pi)) \rightarrow 0$ as $N \rightarrow \infty$, and the Corollary is proved.

Since the functions $\mathbf{r}, \mathbf{r}^+, \mathbf{r}^-$ are positive, rational and homogeneous of degree $-m$, Corollary 7 implies that, for some positive constant $C(\mathcal{R})$, depending only on the Rauzy class $\mathcal{R}$, we have

$$\rho^+(\lambda, \pi) = C(\pi)\mathbf{r}^+(\lambda, \pi), \rho^-(\lambda, \pi) = C(\pi)\mathbf{r}^-(\lambda, \pi).$$

By construction, for any $\lambda \in \mathbb{R}_+^m$ we have

$$\mathbf{r}_\pi^+(\lambda_1, \dots, \lambda_m) = \mathbf{r}_{\pi^{-1}}^-(\lambda_{\pi(1)}, \dots, \lambda_{\pi(m)}).$$

In view of this observation, it suffices to prove only the first assertion of the Lemma 10, as the second one follows automatically.

Take $\delta = (\delta_1, \dots, \delta_m) \in \mathbb{R}^m$, and, for $\theta > 0$, define

$$J_\theta^{(m)}\delta = (\delta_1, \dots, \delta_m + \theta), J_{-\theta}^{(\pi^{-1}m)}\delta = (\delta_1, \dots, \delta_{\pi^{-1}m} - \theta, \dots, \delta_m).$$

Proposition 9 *Let $\theta > 0$. If $\delta \in K_\pi$, then $J_\theta^{(m)}\delta \in K_\pi$, $J_{-\theta}^{(\pi^{-1}m)}\delta \in K_\pi$. If $\delta \in K_\pi^-$, then $J_\theta^{(m)}\delta \in K_\pi^-$. If $\delta \in K_\pi^+$, then $J_{-\theta}^{(\pi^{-1}m)}\delta \in K_\pi^+$.*

This follows directly from the definition of the cones K_π, K_π^-, K_π^+ . From (22) we obtain

$$\begin{aligned} \text{Area}(\lambda, \pi, J_\theta^{(m)}\delta) &= \text{Area}(\lambda, \pi, \delta) + \theta \left(\sum_{r=\pi(m)+1}^m \lambda_{\pi^{-1}r} \right), \\ \text{Area}(\lambda, \pi, J_{-\theta}^{(\pi^{-1}m)}\delta) &= \text{Area}(\lambda, \pi, \delta) + \theta \left(\sum_{r=\pi^{-1}(m)+1}^m \lambda_r \right), \end{aligned}$$

which implies

Proposition 10

$$\text{Area}(\lambda, \pi, \delta) \leq \text{Area}(\lambda, \pi, J_\theta^{(m)}\delta) \leq \text{Area}(\lambda, \pi, \delta) + \theta|\lambda|.$$

$$\text{Area}(\lambda, \pi, \delta) \leq \text{Area}(\lambda, \pi, J_{-\theta}^{(\pi^{-1}m)}\delta) \leq \text{Area}(\lambda, \pi, \delta) + \theta|\lambda|.$$

For $s \in \mathbb{R}$ and a hyperplane of the form $\delta + \dots + \delta_m = s$, let vol_{m-1} stand for the induced $(m-1)$ -dimensional volume form on the hyperplane.

Denote

$$K_{s,\pi} = K_\pi \cap \left\{ \delta : \sum_{i=1}^m \delta_i = s \right\},$$

$$K_s(\lambda, \pi) = K(\lambda, \pi) \cap K_{s,\pi},$$

$$V_s(\lambda, \pi) = \text{vol}_{m-1}(K_s(\lambda, \pi)).$$

Denote by $\mathbf{a}_{\max}^-$ the maximal possible value of $\delta_1 + \dots + \delta_m = -a_m$ in $K(\lambda, \pi)$.

Proposition 11 Assume $0 \leq s \leq \mathbf{a}_{\max}^-$. Then

$$V_s(\lambda, \pi) \leq (1+s)^{m-1} V_0(\lambda, \pi).$$

Proof: Indeed, if $(\lambda, \pi, \delta) \in V_s(\lambda, \pi)$, then Proposition 10 implies

$$(\lambda, \pi, \frac{J_{-s}^{(\pi^{-1}m)}\delta}{1+s}) \in V_0(\lambda, \pi),$$

and the assertion follows.

Proposition 12 Assume s , $0 \leq s \leq 1$ is such that $\frac{s}{1-s} \leq \mathbf{a}_{\max}^-$. Then

$$(\frac{1}{1-s})^{m-1} V_s(\lambda, \pi) \geq V_0(\lambda, \pi).$$

Denote $\theta = \frac{s}{1-s}$, then $s = \frac{\theta}{1+\theta}$. If $(\lambda, \pi, \delta) \in V_0(\lambda, \pi)$, then

$$(\lambda, \pi, \frac{J_{\theta}^{(m)}\delta}{1+\theta}) \in V_s(\lambda, \pi),$$

and, again, the assertion follows.

Propositions 11, 12 imply

Lemma 12 For any $C_1 > 0$ there exists $C_2 > 0$ such that the following is true.

Let $\mathbf{a}_{\max}^-(\lambda, \pi) < C_1$. Then

$$\mathbf{r}^-(\lambda, \pi) < C_2 V_{m-1}^0(\lambda, \pi).$$

Note that there exists $\epsilon > 0$, depending only on $\mathcal{R}$ and such that for any $(\lambda, \pi) \in \Delta(\mathcal{R})$, we have $\mathbf{a}_{\max}^- > \epsilon$. In conjunction with Propositions 11, 12, this implies

Lemma 13 There exists a constant C_3 such that for any $(\lambda, \pi) \in \Delta(\mathcal{R})$, we have

$$\rho^-(\lambda, \pi) \geq C_3 V_0(\lambda, \pi).$$

Since $a_m \leq h_{\pi^{-1}m+1}$, we have

$$\mathbf{a}_{\max}^-(\lambda, \pi) \leq \frac{1}{\lambda_{\pi^{-1}m+1}},$$

which implies the following

Corollary 8 For any $C_4 > 0$ there exists $C_5 > 0$ such that the following is true.

Assume $\lambda_{\pi^{-1}m+1} > C_4$. Then

$$\frac{\mathbf{r}^-(\lambda, \pi)}{\mathbf{r}^+(\lambda, \pi)} < C_5,$$

which implies Lemma 10.

8 Kerckhoff names

In the following two sections, we shall use Kerckhoff's convention of numbering the subintervals of an interval exchange [20]; to avoid confusion, we shall speak of *Kerckhoff names* of subintervals.

Take an interval exchange (λ, π) . A *Kerckhoff naming* on the subintervals of (λ, π) is defined by an arbitrary permutation $i_1, \dots, i_m$ of the symbols $\{1, \dots, m\}$. Once such a permutation is given, we assign names $I_{i_1}, \dots, I_{i_m}$ to the subintervals of (λ, π) , from the left to the right (i.e., the subinterval $[0, \lambda_1)$ is named I_{i_1} , the subinterval $[\lambda_1, \lambda_1 + \lambda_2)$ is named I_{i_2} and so forth).

A Kerckhoff naming of the subintervals of (λ, π) induces a naming on the subintervals of $\mathcal{T}(\lambda, \pi)$ in the following way. Assume $\lambda_m < \lambda_{\pi^{-1}m}$ and the Rauzy operation a was applied to (λ, π) in order to obtain $\mathcal{T}(\lambda, \pi)$. Then the subintervals of $\mathcal{T}(\lambda, \pi)$ are named, from the left to the right, by $I_{i_1}, \dots, I_{i_{\pi^{-1}m}}, I_m, I_{\pi^{-1}m+1}, \dots, I_{m-1}$. If $\lambda_m > \lambda_{\pi^{-1}m}$ and the Rauzy operation b was applied, then the subintervals of $\mathcal{T}(\lambda, \pi)$ are just named, as before, by $I_{i_1}, \dots, I_{i_m}$, from the left to the right. Proceeding inductively, we obtain a naming for any $\mathcal{G}^n(\lambda, \pi)$. Conversely, if we have a Kerckhoff naming of subintervals of (λ, π) , then, for any word $w \in \mathcal{W}_{A,B}$ compatible with (λ, π) , we automatically obtain a Kerckhoff naming on the subintervals of $t_w(\lambda, \pi)$ and $T_w(\lambda, \pi)$.

Let (λ, π) be an interval exchange with a Kerckhoff naming $I_{i_1}, \dots, I_{i_m}$. If $(\lambda, \pi) \in \Delta^+$, then we say that $I_{i_{\pi^{-1}m}}$ is the subinterval *in the critical position* (we shall also sometimes say “in the a -critical position”). If $(\lambda, \pi) \in \Delta^-$, then we say that I_{i_m} is the subinterval *in the critical position* (we shall also sometimes say “in the b -critical position”).

9 Exponential growth.

Let $\mathbf{x} \in \overline{\Delta}$, that is, $\mathbf{x} = (\dots, (\lambda(-n), \pi(-n)), \dots, (\lambda, \pi))$, where, as usual, $\mathcal{G}(\lambda(-n), \pi(-n)) = (\lambda(1-n), \pi(1-n))$. Define the words $w(n)$ by the relation $(\lambda(-n), \pi(-n)) = t_{w(n)}(\lambda, \pi)$. Set $(\Lambda(-n), \pi(-n)) = T_{w(n)}(\lambda, \pi)$.

Lemma 14 *There exists N such that the following is true. For any $\mathbf{x} \in \overline{\Delta}(\mathcal{R})$, there exist $i_1, i_2 \in \{1, \dots, m\}$ such that*

$$\Lambda(-N)_{i_1} + \Lambda(-N)_{i_2} \geq 2(\lambda(0)_{i_1} + \lambda(0)_{i_2})$$

Proof:

Take a point $x \in \overline{\Delta}$,

$$x = (\dots, (\lambda(-n), \pi(-n)), \dots, (\lambda, \pi)).$$

Give Kerckhoff names $I_1, \dots, I_m$ to the subintervals of the exchange (λ, π) from the left to the right, so that the length of I_i is λ_i . We thus automatically obtain a Kerckhoff naming for the subintervals of $((\lambda(-n), \pi(-n)))$ for any n .

Let I_{j_n} be the critical subinterval for $(\lambda(-n), \pi(-n))$.

Consider the infinite sequence

$$I_{j_1} \dots I_{j_n} \dots \quad (23)$$

Note that $j_n \neq j_{n+1}$. A subword $I_{j_k} \dots I_{j_{k+l}}$ will be called a *simple cycle* if $I_{j_k} = I_{j_{k+l}}$ whereas $I_{j_k}, \dots, I_{j_{k+l-1}}$ are all distinct. Naturally, $l \leq m$. There are finitely many possible simple cycles, therefore there exists N , depending only on m , such that for any word of length N in the alphabet $\{I_1, \dots, I_m\}$, some simple cycle occurs at least m times. Now take the word

$$I_{j_1} \dots I_{j_N}, \quad (24)$$

the beginning of the sequence (23), and take a simple cycle which occurs m times, say

$$I_{l_1} \dots I_{l_r}, \quad (25)$$

Here, of course, $r \leq m$. Now estimate the non-renormalized length of the subintervals $I_{l_1}, \dots, I_{l_r}$ ($r \leq m$). In the beginning, these are $\lambda_{l_1}, \dots, \lambda_{l_r}$. The key observation is, as usual, that the interval in critical position at a given inverse Zorich step was, at the previous step, added to the previous critical interval. After the first occurrence of the cycle (25), therefore, the (non-normalized) length of I_{l_1} is at least $\lambda_{l_1} + \lambda_{l_2}$, that of I_{l_2} is at least $\lambda_{l_2} + \lambda_{l_3}$ and so forth. After the second occurrence of (25), the length of I_{l_1} is at least $\lambda_{l_1} + \lambda_{l_2} + \lambda_{l_3}$, that of I_{l_2} is at least $\lambda_{l_2} + \lambda_{l_3} + \lambda_{l_4}$, and so forth. Finally, after the r -th occurrence of (25), the length of I_{l_1} is not less than $\lambda_{l_1} + \lambda_{l_2} + \dots + \lambda_{l_r}$, that is, not less than $2\lambda_{l_1}$, since $\lambda_{l_1} = \lambda_{l_r}$. The Lemma is proven.

10 Proof of the Lemma 5

An informal sketch of the proof of Lemma 5. One divides the subintervals into “big” ones and “small” ones: the aim is to obtain one more “big” interval. For this, one must first put a small subinterval into critical position. This is achieved by Lemma 15. In the previous section, we have seen that the total length of the (non-renormalized) interval grows exponentially with the number of Zorich steps (with an exponent depending on ϵ). When the total length of the interval doubles, we obtain a new “big” subinterval.

10.1 Putting a small interval into critical position

Take an interval exchange (λ, π) and name the subintervals $I_1, \dots, I_m$, from the right to the left.

Proposition 13 *Any interval can be put both in the a -critical and in the b -critical position.*

Proof: First note that if an interval can be put in the a -critical position, then it can also be put into the b -critical position just by performing the entire a -cycle of the corresponding permutation. Since the permutation is irreducible, it suffices to prove that, if I_i can be put into critical position, then also all I_j for $j > i$. To prove this, take the shortest word w that puts I_i into the a -critical position. Then, in the preimage, all I_j , $j > i$, still stand to the right of I_i , though perhaps in a different order (because an inversion of order between I_i and I_j can only happen once I_i reaches the critical position). Therefore, we can immediately place any of the I_j , $j > i$, into the b -critical position, but then also into the a -critical position.

More precisely, pick a positive integer $k \leq m$ and a real $\gamma > 0$. We say that we have a (k, γ) -big-small decomposition if the intervals of the exchange are divided into two groups: $I_{i_1}, \dots, I_{i_k}$, each of length at least γ , and the remaining ones (nothing is said about the length of the remaining ones).

Under the Kerckhoff convention, a big-small decomposition of (λ, π) is inherited by all $t_w(\lambda, \pi)$ (one just takes the intervals with the same names).

Lemma 15 *For any $\gamma > 0$, there exist constants $p(\gamma), L(\gamma)$ such that the following is true. Let $(\lambda, \pi) \in \Delta_{k, \gamma}$ with a fixed big-small decomposition. Then there exists $w \in \mathcal{W}_{A, B}$ such that*

1. $\mathbb{P}(w|\lambda, \pi) \geq p(\gamma)$.
2. $|T_w(\lambda, \pi)| < L(\gamma)$.
3. the exchange $t_w(\lambda, \pi)$ has a small interval in critical position.

Proof: Take the shortest word (in terms of the number of Zorich operations) that puts a small interval into critical position. Among all such words, pick the one that involves the smallest number of Rauzy operations. The length of this word, as well as the number of Rauzy operations involved, only depends on the Rauzy class. At each intermediate Rauzy step, all subintervals following the critical one either in the preimage or in the image must be big, otherwise there would exist a shorter word placing a small interval into critical position. Therefore, by Lemma 9 and the Corollary 6, the probability of each Zorich operation involved is bounded from below by a constant that only depends on γ . The Lemma is proved.

10.2 Completion of the proof.

Proof: Take any $x \in \overline{\Delta}$. Take the first n such that $|\Lambda(-n)| > 2$. By Lemma 14, $n < K|\log \epsilon|$. By Lemma 7, with positive probability depending only on M , we can assume $|\Lambda(-n)| < 2M$. Consider two cases:

1. at all steps from 1 to n , only small intervals were added between themselves.
2. at some step a large interval was added to a small one.

Note, that since we start with a small interval in critical position, either one or the other case holds (for, in order that a small interval be added to a big interval, a big interval must first be placed into critical position, and for that it must first be added to a small one).

In the first case, the lengths of all large intervals remain the same, and after renormalization at step n , each large interval has length at least $\gamma/2M$. However, since $|\Lambda(-n)| > 2$, there must be another interval of length at least $1/2mM$, and the Lemma is proved.

In the second case, let n_1 be the first moment, at which a big interval is added to a small one. Then $|\Lambda(-n_1)| < 2$, and, since at previous moments only small intervals were added between themselves, we have $k+1$ intervals of length at least $\gamma/2$, and the Lemma is proved completely.

11 Return times for the Teichmüller flow.

We have in fact proven a stronger statement, namely, the following Lemma.

Lemma 16 *For any word $\mathbf{q} \in \mathcal{W}_{A,B}$ such that all entries of the matrix $A(\mathbf{q})$ are positive, there exist constants $K_0(\mathbf{q}), p(\mathbf{q})$, depending only on $\mathbf{q}$ and such that the following is true. For any $K \geq K_0$ and any $(\lambda, \pi) \in \Delta(\mathcal{R})$,*

$$\mathbb{P}(\exists n : (\lambda(-n), \pi(-n)) \in \Delta_{\mathbf{q}}, |\Lambda(-n)| < K) | (\lambda, \pi)) \geq p(\mathbf{q})$$

This statement has the following Corollary for the Teichmüller flow on the space of zippered rectangles.

Take an arbitrary word $\mathbf{q} = q_1 \dots q_{2l+1} \in \mathcal{W}_{A,B}$ such that all entries of the matrix $A(q_1 \dots q_l)$ are positive and all entries of the matrix $A(\mathbf{q})$ are positive. As usually, set

$$\Delta_{\mathbf{q}} = \{(\lambda, \pi) : \Phi(\lambda, \pi) = \omega_1 \dots \omega_n \dots, \omega_1 = q_1, \dots, \omega_{2l+1} = q_{2l+1}\}.$$

Consider also the cylinder

$$\overline{\Delta}_{\mathbf{q}} = \{\omega \in \Omega_{A,B}^{\mathbb{Z}}, \omega_{-l} = q_1, \dots, \omega_l = q_{2l+1}\}.$$

Consider the flow P^t as a special flow over $\overline{\Delta}_{\mathbf{q}}$. Denote the roof function of the flow by $\tau_{\mathbf{q}}$.

We shall now see that Lemma 16 implies

Corollary 9 *There exists $\epsilon > 0$ such that*

$$\int_{\overline{\Delta}_{\mathbf{q}}} \exp(\epsilon \tau_{\mathbf{q}}(\omega)) d\mathbb{P}(\omega) < +\infty.$$

Take $\omega \in \Omega_{A,B}^{\mathbb{Z}}$. As usually, set

$$(\lambda(-n), \pi(-n)) = \Phi^{-1}(\omega_{-n} \dots \omega_0 \omega_1 \dots), (\Lambda(-n), \pi(-n)) = T_{\omega_{-n} \dots \omega_{-1} \omega_0}(\lambda(0), \pi(0)).$$

Set $n_{\mathbf{q}}(\omega)$ to be the smallest n such that

$$\omega_{-n} = q_1, \dots, \omega_{-n+2l} = q_{2l+1}.$$

Finally, set $L_{\mathbf{q}}(\omega) = \log |\Lambda(-n_{\mathbf{q}}(\omega))|$. Informally, $L_{\mathbf{q}}(\omega)$ is the “Teichmüller flow time” it takes ω to reach $\bar{\Delta}_{\mathbf{q}}$.

To establish the Corollary 9, it suffices to prove

Proposition 14 *There exists $\epsilon > 0$ such that*

$$\int_{\Omega_{\mathcal{A},B}^{\mathbb{Z}}} \exp(\epsilon L_{\mathbf{q}}(\omega)) d\mathbb{P}(\omega) < +\infty.$$

Proof of Proposition 14. Our main tool will be Lemma 16. Take a $K > K_0$ such that $1 - p(\mathbf{q}) + \frac{1}{K} < 1$. Define a random time $k_1(\omega)$ to be the first moment n such that $|\Lambda(-n)(\omega)| > K$. Note that the map

$$\tilde{\sigma}(\omega) \rightarrow \sigma^{-k_1(\omega)}(\omega)$$

is invertible (here, as always, σ is the shift on $\Omega_{\mathcal{A},B}$).

Introduce a function $\eta : \Omega_{\mathcal{A},B}^{\mathbb{Z}} \rightarrow \mathbb{N}$ by the formula

$$\eta(\omega) = \lceil \frac{\log |\Lambda(-k_1(\omega))|}{\log K} \rceil.$$

In other words, $\eta(\omega) = n$ if

$$K^n \leq |\Lambda(-k_1(\omega))| \leq K^{n+1}.$$

Proposition 15 *There exists a constant C such that the following is true for any $K > K_0$.*

For any $c_1 \dots c_n \dots \in \Omega_{\mathcal{A},B}^+$,

$$\mathbb{P}(\{\omega : \eta(\omega) = n | \omega_1 = c_1, \dots, \omega_n = c_n \dots\}) \leq \frac{C}{K^n}.$$

This immediately follows from Lemma 7.

Proposition 16

$$\mathbb{P}(\{\omega : \eta(\omega) = 1, \omega_{-k_1(\omega)} \dots \omega_0 \text{ does not contain the word } \mathbf{q}\}) \leq 1 - p(\mathbf{q})$$

Finally, take a large N and let

$$n_N(\omega) = \min n : k_1(\omega) + \dots k_1(\tilde{\sigma}^{-n}(\omega)) \leq N.$$

Note that, by definition,

$$K^N \leq |\Lambda(-n_1)(\omega)| \leq K^{2N}.$$

Now consider the set

$$\tilde{\Omega}(N) = \{\omega : \omega_{-n_N(\omega)} \dots \omega_0 \text{ does not contain the word } \mathbf{q}\}.$$

Note that

$$\{\omega : L_{\mathbf{q}}(\omega) > 2N\} \subset \tilde{\Omega}(N).$$

It suffices, therefore, to prove that there exists $r < 1$ such that

$$\mathbb{P}(\tilde{\Omega}(N)) \leq r^N.$$

But by the previous two propositions, we immediately have

$$\mathbb{P}(\tilde{\Omega}) \leq C(1 - p(\mathbf{q}) + \frac{1}{K})^N,$$

and, since $1 - p(\mathbf{q}) + \frac{1}{K} < 1$, the Proposition follows.

This Proposition admits an equivalent formulation in terms of the norms of renormalization matrices on the space of interval exchange transformations.

More precisely, for $(\lambda, \pi) \in \Delta_{\mathbf{q}}$, $\Phi(\lambda, \pi) = \omega_1 \dots \omega_n \dots$, we let $n^{\mathbf{q}}(\lambda, \pi)$ to be the smallest $n > 0$ such that $\mathcal{G}^n(\lambda, \pi) \in \Delta_{\mathbf{q}}$, and we set

$$\mathcal{N}(\lambda, \pi) = \|A(\omega_1 \dots \omega_{n^{\mathbf{q}}(\lambda, \pi)})\|.$$

Corollary 10 *There exists $\epsilon > 0$ such that*

$$\int_{\Delta_{\mathbf{q}}} N(\lambda, \pi)^{\epsilon} d\mathbb{P} < +\infty.$$

Remark. First results on exponential decay for the probabilities of return times were obtained by Jayadev Athreya. In his approach, Athreya used the dynamics of $SL(2, \mathbb{R})$ -action, which allowed him to obtain optimal exponents. The argument above is an attempt to recover some of Athreya's theorems using the language of interval exchange transformations; the argument above does not, however, give an optimal exponent.

Avila, Gouëzel, and Yoccoz have recently announced exponential decay of correlations for the Teichmüller flow. One of the steps in their proof is, again, an exponential estimate for return times, which they have obtained independently (Avila [oral communication]). Their exponent is optimal.

12 Estimate of the measure.

Lemma 17 *There exists a constant $C(\mathcal{R})$ depending only on the Rauzy class $\mathcal{R}$ such that*

$$\nu(\Delta(\mathcal{R}) \setminus \Delta_{\epsilon}(\mathcal{R})) < C\epsilon$$

The proof repeats that of Proposition 13.2 in Veech [1].
 Lemma 4 and Corollary 5 therefore imply the following

Corollary 11 *Let $\mathbf{q} \in W_{A,B}$, $\mathbf{q} = q_1 \dots q_l$ be such that all entries of the matrix $A(\mathbf{q})$ are positive.*

There exist $C > 0, \alpha > 0$ such that the following is true for any n .

$$\mathbb{P}((\lambda, \pi) : \mathcal{G}^{2k}(\lambda, \pi) \notin \Delta(\mathbf{q}) \text{ for all } k, 1 \leq k \leq n) \leq C \exp(-\alpha\sqrt{n}).$$

Proof: Let $n = r^2$ and denote

$$X(n, \mathbf{q}) = \{(\lambda, \pi) : \mathcal{G}^{2k}(\lambda, \pi) \notin \Delta(\mathbf{q}) \text{ for all } k, 1 \leq k \leq n\}.$$

Take

$$B(n) = \{(\lambda, \pi) : \mathcal{G}^{2k}(\lambda, \pi) \notin \Delta_{\exp(-r)} \text{ for some } k, 1 \leq k \leq n\}$$

Then, by the previous Lemma, $\nu(B(n)) \leq Cr^2 \exp(-r)$, whereas, by Corollary 5,

$$\nu(X(n, \mathbf{q}) \setminus B(n)) \leq (1 - p(\mathbf{q}))^r,$$

and Corollary 11 is proven.

Remark. This result allows to use the tower method of L.-S. Young [11] and to obtain the decay rate $\exp(-\alpha\sqrt{n})$ for correlations of bounded Hölder functions. For bounded Lipschitz functions, one can also use the method of V. Maume-Deschamps [12] and obtain the uniform rate of decay at the rate $\exp(-\alpha n^{1/2-\epsilon})$. It is not clear to me, however, how to use either of these methods in the invertible case.

13 Inequalities

Let

$$W_{A,B}^+ = \{w \in W_{A,B} : |w| \text{ is even, } \Delta(w) \subset \Delta^+\}.$$

Lemma 18 *For any $C_1, C_2 > 0$ there exists $C_3 > 0$ such that the following is true.*

Suppose $\text{row}(A) < C_1$ and $\lambda \in \Delta_{C_2}$.

Then

$$\frac{1}{C_3} \leq \frac{|A\lambda|^m}{\prod_{j=1}^m \sum_{i=1}^m A_{ij}} \leq C_3$$

Proof:

Denote $A_j = \sum_{i=1}^m A_{ij}$, so that $|A| = \sum_{j=1}^m A_j$.

Then

$$\frac{A_j}{A_k} \leq \text{row}(A),$$

whence

$$\frac{A_j}{|A|} \geq \frac{1}{m \operatorname{row}(A)}.$$

Finally, if $\lambda \in \Delta_{C_2}$, then

$$|A\lambda| \geq C_2|A|,$$

which completes the proof.

Corollary 12 *For any $C_4 > 0$, $C_5 > 0$ there exists $C_6 > 0$ such that the following is true. Suppose $(\lambda, \pi) \in \Delta_{C_4}$. Suppose $w \in \mathcal{W}_{\mathcal{A},B}$ is compatible with (λ, π) and such that $\operatorname{row}(A(w)) < C_5$. Then*

$$\frac{1}{C_6} \leq \frac{\mathbf{m}(C(w))}{\mathbb{P}(w|(\lambda, \pi))} \leq C_6$$

Corollary 13 *For any $C_7 > 0$, $C_8 > 0$ $C_9 > 0$, there exists $C_{10} > 0$ such that the following is true.*

Suppose $(\lambda, \pi) \in \Delta_{C_7}$.

Suppose $w \in \mathcal{W}_{\mathcal{A},B}$ is compatible with (λ, π) and furthermore satisfies

$$\operatorname{row}(A(w)) < C_8, \quad \Delta(w) \subset \Delta_{C_9}$$

Then

$$\frac{1}{C_{10}} \leq \frac{\mathbb{P}(C(w))}{\mathbb{P}(w|(\lambda, \pi))} \leq C_{10}$$

Corollary 14 *Let M be such that for any $n > M$ any two vertices in the Rauzy graph can be joined in n steps.*

Then for any $C_{17} > 0$, $C_{18} > 0$ $C_{19} > 0$, there exists $C_{20} > 0$ such that the following is true.

Suppose $(\lambda, \pi) \in \Delta^+ \cap \Delta_{C_{17}}$.

Suppose $w \in W_{\mathcal{A},B}^+$ satisfies

$$\operatorname{row}(A(w)) < C_{18}, \quad \Delta(w) \subset \Delta^+ \cap \Delta_{C_{19}}$$

Then for any $n \geq M$, we have

$$\frac{1}{C_{20}} \leq \frac{\mathbb{P}(C(w))}{\mathbb{P}^{(2n)}(w|(\lambda, \pi))} \leq C_{20}$$

From the definition (4) of the Hilbert metric it easily follows that for any $\lambda, \lambda' \in \Delta_{m-1}$ we have

$$e^{-d(\lambda, \lambda')} \lambda'_i \leq \lambda_i \leq e^{d(\lambda, \lambda')} \lambda'_i. \quad (26)$$

Proposition 17 *Assume $\lambda, \lambda' \in \Delta_\pi^+$. Then*

$$\exp(-md(\lambda, \lambda')) \leq \frac{\rho(\lambda, \pi)}{\rho(\lambda', \pi)} \leq \exp(md(\lambda, \lambda'))$$

Proof. Indeed, there exist linear forms

$$l_i^{(j)}(\lambda) = \sum_{k=1}^m a_{ik}^{(j)} \lambda_k,$$

where $a_{ik}^{(j)}$ are nonnegative integers (in fact, either 0 or 1, but we do not need this here),
such that

$$\rho(\lambda, \pi) = \sum_{j=1}^s \frac{1}{l_1^{(j)}(\lambda) l_2^{(j)}(\lambda) \dots l_m^{(j)}(\lambda)}.$$

Clearly, if for all $i = 1, \dots, m$ and some $\alpha > 0$, we have $\alpha^{-1} \lambda_i \leq \lambda'_i \leq \alpha \lambda_i$, then

$$\alpha^{-m} \leq \frac{\rho(\lambda, \pi)}{\rho(\lambda', \pi)} \leq \alpha^m,$$

and the Proposition is proved.

For similar reasons we have

Proposition 18 *Assume $\lambda, \lambda' \in \Delta_\pi^+$ and let A be an arbitrary matrix with nonnegative integer entries. Then*

$$\exp(-md(\lambda, \lambda')) \leq \frac{\rho(A\lambda, \pi)}{\rho(A\lambda', \pi)} \leq \exp(md(\lambda, \lambda'))$$

From these propositions and the formula 11 we obtain

Corollary 15 *Let $c \in \mathcal{A}$ be compatible with π . Then for any $\lambda, \lambda' \in \Delta_\pi^+$ we have*

$$\exp(-2md(\lambda, \lambda')) \leq \frac{\mathbb{P}(c|(\lambda, \pi))}{\mathbb{P}(c|(\lambda', \pi))} \leq \exp(2md(\lambda, \lambda'))$$

This Corollary implies the following

Lemma 19 *Let $w \in W_{\mathcal{A}, B}^+$ be such that the cylinder $C(w)$ has finite Hilbert diameter.*

Then for any c compatible with w and any $(\lambda_0, \pi) \in C(w)$ we have

$$\exp(-2m \operatorname{diam} C(w)) \leq \frac{\mathbb{P}(c|(\lambda_0, \pi))}{\mathbb{P}(\omega_0 = c|\omega|_{[1, |w|]} = w)} \leq \exp(2m \operatorname{diam} C(w))$$

Proof: We have

$$\nu(C(cw)) = \int_{C(w)} \mathbb{P}(c|(\lambda, \pi)) d\nu(\lambda, \pi)$$

Let $d = \text{diam}C(w)$. For any $(\lambda, \pi), (\lambda', \pi) \in C(w)$, we have, by Corollary 15,

$$\exp(-2md) \leq \frac{\mathbb{P}(c|(\lambda, \pi))}{\mathbb{P}(c|(\lambda', \pi))} \leq \exp(2md).$$

Fix an arbitrary $(\lambda_0, \pi) \in \Delta_w$.

Then, from the above,

$$\begin{aligned} \nu(C(w))P(c|(\lambda_0, \pi))\exp(-2md) &\leq \int_{C(w)} P(c|(\lambda, \pi))d\nu(\lambda, \pi) \leq \\ &\leq \nu(C(w))P(c|(\lambda_0, \pi))\exp(2md), \end{aligned}$$

and, since, by definition, we have

$$\mathbb{P}(\omega_0 = c|\omega|_{[1, |w|]} = w) = \frac{\mathbb{P}(cw)}{\mathbb{P}(w)},$$

the Lemma is proved.

For $N \in \mathbb{N}$ and $A \subset \Delta(\mathcal{R})$, we denote $\mathbb{P}^{(N)}(A|(\lambda, \pi)) = \mathbb{P}((\lambda(-N), \pi(-N)) \in A | (\lambda(0), \pi(0)) = (\lambda, \pi))$; for $w \in \mathcal{W}_{A,B}$, we write $\mathbb{P}^{(N)}(w|(\lambda, \pi)) = \mathbb{P}^{(N)}(\Delta(w)|(\lambda, \pi))$.

Lemma 20 *Let M be a number such that for any $N \geq M$ any two vertices of the Rauzy graph can be connected in N steps. For any $\gamma > 0$, $N \geq M$ there exists a constant C_0 depending only on γ and N such that for any word $w \in \mathcal{W}_{A,B}^+$ and any $(\lambda, \pi) \in \Delta_\gamma$*

$$\mathbb{P}^{(2N)}(w|(\lambda, \pi)) \geq \frac{C_0}{|A(w)\lambda|^m}$$

Proof:

Let $w = w_1 \dots w_{2n}$, and let $w_{2n} = (a, m_1, \pi_1)$.

Let $\pi'_1 \pi'_2 \dots \pi'_{2N}$ a path of length $2N$ between π and π_1 (here $\pi'_1 = \pi$, $\pi'_{2n} = \pi_1$, $\pi_{2k+1} = a\pi_{2k}$, $\pi_{2k+2} = b\pi_{2k+1}$).

Denote $w_{n+2i+1} = (a, 1, \pi_{2i+1})$, $w_{n+2i} = (b, 1, \pi_{2i})$. In other words, the word $= w_{2n+1} \dots w_{2n+2N} \in \mathcal{W}_{A,B}$ is the word corresponding to the path $\pi'_1 \pi'_2 \dots \pi'_{2N}$ in the Rauzy graph. Then $w' = w_1 \dots w_{2n+2N}$ is a word compatible with (λ, π) . Besides,

$$|A(w_{2n+1}c_{n+2} \dots w_{2n+2N})| < (2N)^{(2N)}.$$

We have

$$P^{(2n)}(w|(\lambda, \pi)) \geq P(w'|(\lambda, \pi)) = \frac{\rho(T_{w'}(\lambda), w'\pi)}{|A(w')\lambda|^m \rho(\lambda, \pi)},$$

There exists a universal constant C_1 such that $\rho(\lambda', \pi') > C_1$ for any $(\lambda', \pi') \in \Delta^+$ (the density of the invariant measure is bounded from below).

Then, $|A(w')\lambda|^m \leq |A(w')|^m \leq (2N)^{2mN} |A(w)|^m$.

Finally, there exists a C_2 depending on c only such that if $\lambda_i > c$ for all i then $\rho(\lambda, \pi) > C_2$.

Combining all of the above, we obtain the result of the Lemma.

14 Markov approximation and the Doeblin condition

14.1 Good cylinders

Let $\mathbf{q} = q_1 \dots q_l$ be a word such that all entries of the matrix $A(\mathbf{q})$ are positive. Fix $\epsilon > 0$ and let k_0 be such that

$$\mathbb{P}(\Delta(\mathbf{q}) \cap \mathcal{G}^{-2n} \Delta(\mathbf{q})) \geq \epsilon \text{ for } n > k_0. \quad (27)$$

Note that, due to mixing, Corollary 5 implies the following

Proposition 19 *Let $\mathbf{q} \in W_{A,B}$, $\mathbf{q} = q_1 \dots q_l$ be such that all entries of the matrix $A(\mathbf{q})$ are positive and that $\Delta(\mathbf{q}) \subset \Delta^+$. Then there exist positive constants $K(\mathbf{q}), p(\mathbf{q})$ such that the following is true for any $\epsilon > 0$. Suppose $(\lambda, \pi) \in \Delta_\epsilon \cap \Delta^+$ and set n to be the integer part of $K(\mathbf{q})|\log \epsilon|$. Then*

$$\mathbb{P}\{(\lambda(-2n), \pi(-2n)) \in \Delta(\mathbf{q}) | (\lambda(0), \pi(0)) = (\lambda, \pi)\} \geq p(\mathbf{q}).$$

Take $k \geq k_0$. Let $r = 2(K+1)k + 2M$, where K is the constant from the Lemma 4 and M is the connecting constant of the Rauzy graph from Lemma 20.

Let θ , $0 < \theta < 1$ be arbitrary. A word $w = w_1 \dots w_k$ is called *good* if

1. $\Delta(w) \subset \Delta_{\exp(-k)}$.
2. the word $\mathbf{q}$ appears at least $\frac{k^\theta}{l}$ times in w (we only count disjoint appearances).

A word $w_1 \dots w_r$ is called good if $w_1 \dots w_k$ is good, a word $w_1 \dots w_{Nr}$ is called good if all words $w_1 \dots w_r$, $w_{r+1} \dots w_{2r}$, $\dots$, $w_{(N-1)r+1} \dots w_{Nr}$ are good, and a word $w_1 \dots w_{Nr+L}$, $L < r$, is good if $w_1 \dots w_{Nr}$ is good and either $L < k$ or $w_{Nr+1} \dots w_{Nr+k}$ is good.

We denote by $\mathbf{G}(N)$ the set of all good words of length N .

Let

$$\Delta(\mathbf{G}(N)) = \cup_{w \in \mathbf{G}(N)} \Delta(w),$$

and

$$\Delta(B(N)) = \Delta^+ \setminus \Delta(\mathbf{G}(N))$$

By Corollary 11, there exist constants C_{31}, C_{32} such that for all r we have

$$\mathbb{P}(\Delta(B(N)) \leq C_{31}N \exp(-C_{32}r^{(1-\theta)/2}). \quad (28)$$

and, for any $(\lambda, \pi) \in \Delta(\mathbf{q})$, also

$$\mathbb{P}((\lambda(-1), \pi(-1)) \in \Delta(B(N)) | (\lambda(0), \pi(0)) = (\lambda, \pi)) \leq C_{31}N \exp(-C_{32}r^{(1-\theta)/2}). \quad (29)$$

14.2 Preliminary estimates for the Doeblin condition.

From Corollary 15 we deduce that there exists a constant C_{33} such that for any $(\lambda, \pi), (\lambda', \pi) \in \Delta(\mathbf{q})$, and any word w compatible with $\mathbf{q}$, we have

$$\frac{1}{C_{33}} \leq \frac{\mathbb{P}(w|(\lambda, \pi))}{\mathbb{P}(w|(\lambda', \pi))} \leq C_{33}.$$

Finally, by Lemma 20, there exists a constant C_{34} such that for any $w \in \mathcal{W}_{A,B}$ and for any $N > M$ we have

$$\frac{1}{C_{34}} \leq \frac{\mathbb{P}^{(2N)}(w|(\lambda, \pi))}{\mathbb{P}^{(2N)}(w|(\lambda', \pi))} \leq C_{34}.$$

Take an arbitrary point $(\lambda, \pi) \in \Delta_{\mathbf{q}}$. Define a new measure φ on Δ^+ . Namely, for a set $A \subset \Delta^+$ put

$$\varphi(A) = \mathbb{P}(\lambda(-2M), \pi(-2M)) \in A | \lambda(0), \pi(0) = (\lambda, \pi)) \quad (30)$$

Lemma 21 *There exists a constant $\alpha > 0$ such that the following is true for any r . Let $\mathcal{C}_1, \mathcal{C}_2 \in \mathbf{G}(r)$.*

Then

$$\mathbb{P}(\omega|_{[1,r]} = \mathcal{C}_1, \omega|_{[r+1,2r]} \in \mathbf{G}(r) | \omega|_{[2r+1,3r]} = \mathcal{C}_2) \geq \alpha \varphi(\mathcal{C}_1)$$

Indeed, we have the following propositions:

Proposition 20 *There exist a constant p_1 such that the following is true for all r and all $n \geq r$.*

Let $\mathcal{C}_2 \in \mathbf{G}(r)$, $(\lambda, \pi) \in \mathcal{C}_2$. Then

$$\mathbb{P}((\lambda(-2n), \pi(-2n)) \in \Delta(\mathbf{q}) | (\lambda(0), \pi(0)) = (\lambda, \pi)) \geq p_1.$$

This follows from the definition of a good cylinder and Corollary 5.

Proposition 21 *There exists a constant p_2 such that the following is true for all k .*

$$\mathbb{P}(\omega|_{[1,r]} \in \mathbf{G}(r), \omega|_{[2M+1,l+2M+1]} = \mathbf{q} | \omega|_{[r+1,r+l+1]} = \mathbf{q}) \geq p_2$$

This follows from the estimates (28),(29) on the measure of bad cylinders and from Proposition 19.

Proposition 22 *There exists a constant p_3 such that the following is true for all r . Let $c_1 \dots c_n \dots \in \Delta(\mathbf{q})$.*

$$\mathbb{P}(\omega|_{[1,r]} = \mathcal{C}_1 | \omega_{r+2M+1} = c_1, \omega_{r+2M+2} = c_2, \dots) \geq p_3 \varphi(\mathcal{C}_1)$$

This follows directly from Lemma 20.

The three Propositions imply Lemma 21.

14.3 Approximation by a Markov measure

We define a new measure $\mathbf{p}_{r,\theta}$ on the set $\mathbf{G}(r^2)$ of good cylinders of length r^2 .

Let $\mathcal{C} = c_1 \dots c_{r^2}$ be a (r, θ) -good cylinder. Set $\mathcal{C}_i = c_{ir+1} \dots c_{(i+1)r}$.

Define

$$\mathbf{p}_{r,\theta}(\mathcal{C}) = \mathbb{P}(\omega|_{[1,r]} = \mathcal{C}_1 | \omega|_{[r+1,2r]} = \mathcal{C}_2) \mathbb{P}(\omega|_{[r+1,2r]} = \mathcal{C}_2 | \omega|_{[2r+1,3r]} = \mathcal{C}_3) \dots \mathbb{P}(\omega|_{[r^2-r+1,r^2]} = \mathcal{C}_r).$$

If D is not a good cylinder, then $\mathbf{p}_{r,\theta}(D) = 0$.

Normalize to get a probability measure:

$$\mathbf{P}_{r,\theta}(\mathcal{C}) = \frac{\mathbf{p}_{r,\theta}(\mathcal{C})}{\sum_{\mathcal{D} \in \mathbf{G}(r^2)} \mathbf{p}_{r,\theta}(\mathcal{D})}.$$

$\mathbf{P}_{r,\theta}$ is a Markov measure of memory r (in general, non-homogeneous), as is shown by the following well-known Lemma [14].

Lemma 22 *For any k , $0 < k < r$, we have*

$$\mathbf{P}_{r,\theta}(\omega|_{[kr+1,(k+1)r]} = \mathcal{C}_k | \omega|_{[(k+1)r+1,r^2]} = \mathcal{C}_{k+1} \dots \mathcal{C}_r) =$$

$$\mathbf{P}_{r,\theta}(\omega|_{[kr+1,(k+1)r]} = \mathcal{C}_k | \omega|_{[(k+1)r+1,(k+2)r]} = \mathcal{C}_{k+1}).$$

From the Hölder property for the transition probability, we have

Proposition 23 *There exist constants C_{41}, C_{42} such that the following is true for any r .*

Let $c_1 \dots c_n \dots \in \Omega_{A,B}$ and assume $c_{n+1} \dots c_{n+r} \in \mathbf{G}(r)$. Then

$$\begin{aligned} & \exp(-C_{41} \exp(-C_{42} k^\theta)) \leq \\ & \leq \frac{P(\omega_1 = c_1, \dots, \omega_n = c_n | \omega_{n+1} = c_{n+1}, \dots, \omega_{n+r} = c_{n+r})}{\mathbb{P}(\omega_1 = c_1, \dots, \omega_n = c_n | \omega_{n+1} = c_{n+1}, \dots, \omega_{n+i} = c_{n+i}, \dots)} \leq \\ & \leq \exp(C_{41} \exp(-C_{42} k^\theta)) \end{aligned}$$

Corollary 16 *There exist constants C_{43}, C_{44} such that the following is true for any r . Let $A \in \mathcal{F}_n$, let $c_{n+1} \dots c_{n+i} \dots \in \Omega_A$, and assume $c_{n+1} \dots c_{n+r} \in \mathbf{G}(r)$. Then*

$$\exp(-C_{43} \exp(-C_{44} k^\theta)) \leq \frac{\mathbb{P}(A | \omega_{n+1} = c_{n+1}, \dots, \omega_{n+r} = c_{n+r})}{\mathbb{P}(A | \omega_{n+1} = c_{n+1}, \dots, \omega_{n+i} = c_{n+i}, \dots)} \leq \exp(C_{43} \exp(-C_{44} k^\theta))$$

Applying l times, we obtain

Lemma 23 *There exist constants $C_{45}, C_{46}, C_{47}, C_{48}$ such that the following is true for any r . Let $c_1 \dots c_{r^2} \in \mathbf{G}(r^2)$. Then for any l , $1 \leq l \leq r$, we have*

$$\exp(-C_{45} l \exp(-C_{46} k^\theta)) \leq$$

$$\begin{aligned} &\leq \frac{\mathbb{P}(\omega_1 = c_1, \dots, \omega_{lr} = c_{lr} | \omega_{lr+1} = c_{lr+1}, \dots, \omega_{r^2} = c_{r^2})}{\mathbf{p}_{r,\theta}(\omega_1 = c_1, \dots, \omega_{lr} = c_{lr} | \omega_{lr+1} = c_{lr+1}, \dots, \omega_{r^2} = c_{r^2})} \leq \\ &\leq \exp(C_{45}l \exp(-C_{46}k^\theta)) \end{aligned}$$

and

$$\exp(-C_{47}l \exp(-C_{48}k^\theta)) \leq \frac{\mathbb{P}(\omega_1 = c_1, \dots, \omega_{lr} = c_{lr})}{\mathbf{p}_{r,\theta}(\omega_1 = c_1, \dots, \omega_{lr} = c_{lr})} \leq \exp(C_{47}l \exp(-C_{48}k^\theta))$$

Summing over cylinders of length lr , we obtain

Corollary 17 *There exist constants C_{49}, C_{50} such that the following is true for any r . Let $c_1 \dots c_{r^2} \in \mathbf{G}(r^2)$. Then for any l , $1 \leq l \leq r$, and any $A \in \mathcal{F}_{lr}$, we have*

$$\exp(-C_{49}l \exp(-C_{50}k^\theta)) \leq \frac{\mathbb{P}(A \cap \mathbf{G}(lr) | \omega_{lr+1} = c_{lr+1}, \dots, \omega_{r^2} = c_{r^2})}{\mathbf{p}_{r,\theta}(A | \omega_{lr+1} = c_{lr+1}, \dots, \omega_{r^2} = c_{r^2})} \leq \exp(C_{49}l \exp(-C_{50}k^\theta))$$

and

$$\exp(-C_{49}l \exp(-C_{50}k^\theta)) \leq \frac{\mathbb{P}(A \cap \mathbf{G}(lr))}{\mathbf{p}_{r,\theta}(A)} \leq \exp(C_{49}l \exp(-C_{50}k^\theta))$$

Using (28), we can estimate the total mass of the measure $\mathbf{p}_{r,\theta}$.

Corollary 18 *There exist constants C_{51}, C_{52} such that for any r we have*

$$\mathbf{p}_{r,\theta}(\mathbf{G}(r^2)) \geq \exp(-C_{51}r \exp(-C_{52}k^{(1-\theta)/2}))$$

We now have normalized versions of previous statements.

Corollary 19 *There exist constants $C_{53}, C_{54}, C_{55}, C_{56}$ such that the following is true for any r . Let $c_1 \dots c_{r^2} \in \mathbf{G}(r^2)$. Then for any l , $1 \leq l \leq r$, and any $A \in \mathcal{F}_{lr}$, we have*

$$\begin{aligned} &\exp(-C_{53}l \exp(-C_{54}k^\theta) - C_{55}r \exp(-C_{56}k^{(1-\theta)/2})) \leq \\ &\leq \frac{\mathbb{P}(A \cap \mathbf{G}(lr) | \omega_{lr+1} = c_{lr+1}, \dots, \omega_{r^2} = c_{r^2})}{\mathbf{p}_{r,\theta}(A | \omega_{lr+1} = c_{lr+1}, \dots, \omega_{r^2} = c_{r^2})} \leq \\ &\leq \exp(C_{53}l \exp(-C_{54}k^\theta) + C_{55}r \exp(-C_{56}k^{(1-\theta)/2})) \end{aligned}$$

and

$$\begin{aligned} &\exp(-C_{53}l \exp(-C_{54}k^\theta) - C_{55}r \exp(-C_{56}k^{(1-\theta)/2})) \leq \\ &\leq \frac{\mathbb{P}(A \cap \mathbf{G}(lr))}{\mathbf{p}_{r,\theta}(A)} \leq \\ &\leq \exp(C_{53}l \exp(-C_{54}k^\theta) + C_{55}r \exp(-C_{56}k^{(1-\theta)/2})). \end{aligned}$$

Using the Markov approximation, we can estimate conditional measure of good cylinders for the measure $\mathbb{P}$:

Corollary 20 *There exist constants $C_{57}, C_{58}, C_{59}, C_{60}$ such that the following is true for any r . Let $c_1 \dots c_{r^2} \in \mathbf{G}(r^2)$. Then for any l , $1 \leq l \leq r$, we have*

$$\mathbb{P}((\omega_1 \dots \omega_{lr}) \in \mathbf{G}(lr) | \omega_{lr+1} = c_{lr+1}, \dots, \omega_{r^2} = c_{r^2}) \geq \exp(-C_{57}l \exp(-C_{58}k^\theta) - C_{59}r \exp(-C_{60}k^{(1-\theta)/2}))$$

Proof: Indeed,

$$\mathbf{P}_{r,\theta}((\omega_1 \dots \omega_{lr}) \in \mathbf{G}(lr) | \omega_{lr+1} = c_{lr+1}, \dots, \omega_{r^2} = c_{r^2}) = 1.$$

14.4 Doeblin Condition

Proposition 24 *There exists C_{61} such that the following holds for any r . For any $C_1 \subset \Delta(\mathbf{q})$, $C_2 \subset \Delta_{\mathbf{q}}$, and any $C_3 \in \mathbf{G}(r)$, we have either*

$$\frac{1}{C_{61}} \leq \frac{\mathbf{p}_{r,\theta}(C_3|C_2)}{\mathbf{p}_{r,\theta}(C_3|C_1)} \leq C_{61},$$

$$\text{or } \mathbf{p}_{r,\theta}(C_3|C_2) = \mathbf{p}_{r,\theta}(C_3|C_1) = 0.$$

Considering n -step transition probabilities, we obtain

Proposition 25 *There exists a constant C_{62} such that the following holds for any r . For any $C_1 \subset \Delta(\mathbf{q})$, $C_2 \subset \Delta_{\mathbf{q}}$ any $C_3 \in \mathbf{G}(r)$, and any $n \geq M$, we have*

$$\frac{1}{C_{62}} \leq \frac{\mathbf{p}_{r,\theta}(\omega|_{[1,r]} = C_1 | \omega_{[2n+r, 2n+2r]} = C_2)}{\mathbf{p}_{r,\theta}(\omega|_{[1,r]} = C_1 | \omega_{[2n+r, 2n+2r]} = C_3)} \leq C_{62}.$$

Now, mixing, Proposition 19 and Proposition 20, and the definition of a good cylinder imply that

Proposition 26 *There exists a constant C_{63} such that the following holds for any r . For any $C_1, C_2, C_3 \in \mathbf{G}(r)$ we have*

$$\frac{1}{C_{63}} \leq \frac{\mathbf{p}_{r,\theta}(\omega|_{[1,r]} = C_1 | \omega_{[2r, 3r]} = C_2)}{\mathbf{p}_{r,\theta}(\omega|_{[1,r]} = C_1 | \omega_{[2r, 3r]} = C_3)} \leq C_{63}.$$

Now let $c_1 \dots c_{r^2} \in \mathbf{G}(r^2)$. Denote $\mathcal{C}_i = c_{ir+1} \dots c_{(i+1)r}$. Lemma 21, together with the above estimates, implies the following

Corollary 21 *There exist constants C_{71}, C_{72} such that the following is true. For any l , $1 \leq l \leq r$, we have*

$$\mathbb{P}(\omega|_{[1,lr]} \in \mathbf{G}(lr), \omega|_{[lr+1, (l+1)r]} = \mathcal{C}_l, \omega|_{[(l+1)r+1, (l+2)r]} \in \mathbf{G}(r) | \omega_{[(l+2)r+1, (l+3)r]} = \mathcal{C}_3) \geq C_{71} \times \varphi(\mathcal{C}_l)$$

and

$$\mathbf{P}_{r,\theta}(\omega|_{[1,lr]} \in \mathbf{G}(lr), \omega|_{[lr+1, (l+1)r]} = \mathcal{C}_l, \omega|_{[(l+1)r+1, (l+2)r]} \in \mathbf{G}(r) | \omega_{[(l+2)r+1, (l+3)r]} = \mathcal{C}_3) \geq C_{72} \times \varphi(\mathcal{C}_l)$$

This is the Doeblin Condition for the measure $\mathbf{P}_{r,\theta}$ (see [13], [14], [22]). The Doeblin Condition implies that there exist constants C_{73}, C_{74} such that for any $\mathcal{C}_1, \mathcal{C}_2 \in \mathbf{G}(r)$, we have

$$\exp(-C_{73} \exp(-C_{74}r)) \leq \frac{\mathbf{P}_{r,\theta}(\omega|_{[1,r]} = \mathcal{C}_1 | \omega|_{[r^2, r^2+r]} = \mathcal{C}_2)}{\mathbf{P}_{r,\theta}(\mathcal{C}_1)} \leq \exp(C_{73} \exp(-C_{74}r)),$$

whence we obtain

Proposition 27 *There exist constants $C_{81}, C_{82}, C_{83}, C_{84}$ such that the following is true for any r .*

$$\begin{aligned} & \exp(-C_{81}(\exp(-C_{82}r) + \exp(-C_{83}r^\theta) + \exp(-C_{84}r^{(1-\theta)/2}))) \leq \\ & \leq \frac{\mathbb{P}(\omega|_{[1,r]} = \mathcal{C}_1 | \omega|_{[r+1, r^2]} \in \mathbf{G}(r^2 - r), \omega|_{[r^2, r^2+r]} = \mathcal{C}_2)}{\mathbb{P}(\mathcal{C}_1)} \leq \\ & \leq \exp(C_{81} \exp(-C_{82}r) + \exp(-C_{83}r^\theta) + \exp(-C_{84}r^{(1-\theta)/2}))). \end{aligned}$$

Moreover, in view of mixing, Proposition 19, and Proposition 20, the same estimate, upto a constant, takes place for any $n \geq r^2$.

Proposition 28 *There exist constants $C_{85}, C_{86}, C_{87}, C_{88}$ such that the following is true for all r and all $n \geq r^2$.*

$$\begin{aligned} & \exp(-C_{85}(\exp(-C_{86}r) + \exp(-C_{87}r^\theta) + \exp(-C_{88}r^{(1-\theta)/2})))) \leq \\ & \leq \frac{\mathbb{P}(\omega|_{[1,r]} = \mathcal{C}_1 | \omega|_{[r+1, n]} \in \mathbf{G}(n - r), \omega|_{[n, n+r]} = \mathcal{C}_2)}{\mathbb{P}(\mathcal{C}_1)} \leq \\ & \leq \exp(C_{85}(\exp(-C_{86}r) + \exp(-C_{87}r^\theta) + \exp(-C_{88}r^{(1-\theta)/2}))))). \end{aligned}$$

15 Approximation of Hölder Functions and Completion of the Proof of Theorems 4, 7, 8.

We shall prove the decay of correlations for a slightly more general class of functions on $\Delta(\mathcal{R})$ than Hölder functions. (we shall need this slightly more general class in the proof of the Central Limit Theorem).

Namely, we shall only require that a function be Hölder in restriction to cylinders of some given length and we shall also allow a moderate growth of the Hölder constant at infinity.

Formally, say that a function $\phi : \Delta(\mathcal{R}) \rightarrow \mathbb{R}$ is *weakly l, α -Hölder* if the following holds. Let k be a positive integer, and let $w \in \mathcal{W}_{\mathcal{A}, B}$, $|w| \leq l$ be such that $\Delta(w) \subset \Delta_{\exp(-k)}$. Then there exists a constant $C(\phi)$ such that for any $(\lambda, \pi), (\lambda', \pi) \in \Delta(w)$, we have

$$|\phi(\lambda, \pi) - \phi(\lambda', \pi)| \leq Ckd(\lambda, \lambda')^\alpha.$$

The smallest such C for a given ϕ will be denoted $C_{l,\alpha}^{weak}(\phi)$. Clearly, if ϕ is Hölder with exponent α , then it is also weakly l, α -Hölder for any l and $C_{l,\alpha}^{weak}(\phi) \leq C_\alpha(\phi)$.

Recall that $\mathcal{B}_n$ is the σ -algebra of sets of the form $\mathcal{G}^{-n}(A)$, $A \subset \Delta(\mathcal{R})$.

To prove the decay of correlations, it suffices to estimate the L_2 -norm of $E(\phi|\mathcal{B}_{2n})$ for a given weakly l, α -Hölder ϕ .

It will be convenient to assume that $\phi \geq 1$ (by linearity, it suffices to consider that case).

Proposition 29 *Let $\theta \in \mathbb{R}$, $0 < \theta < 1$. Let $p > 2$ and $\alpha > 0$. There exist constants C_{91}, C_{92}, C_{93} such that the following is true for any r and any $n \geq r^2$.*

Let $l \leq r$. Let $\phi \in L_p(\Delta(\mathcal{R})^+, \nu)$ be weakly l, α -Hölder and satisfy $\phi \geq 1$.

Then $\phi = \phi_1 + \phi_2 + \phi_3$ where

1. $\phi_1 \geq 1$ on $\mathbf{G}(n)$ and $\phi_1 = \phi_2 = 0$ on $\Delta(B(n))$.

2. for any $(\lambda, \pi) \in \mathbf{G}(n)$, we have

$$\left| \frac{E(\phi_1|\mathcal{F}_n)(\lambda, \pi)}{E(\phi_1)} - 1 \right| \leq \exp(-C_{91}(r^{(1-\theta)/2} + r^\theta)).$$

3. for $(\lambda, \pi) \in G(n)$, we have $|\phi_2| \leq C_{l,\alpha}^{weak}(\phi) \exp(-C_{92}r^\theta)$.

4. $\|\phi_3\|_{L_2} \leq \exp(-C_{93}r^{(1-\theta)/2})\|\phi\|_{L_p}$.

Proof: For any good word $w = w_1 \dots w_{n+r}$, consider its beginning $w_1 \dots w_r$ and choose a point $x_{w_1 \dots w_r} \in \Delta(w_1 \dots w_r)$.

Denote by $\chi_{\Delta(w)}$ the characteristic function of $\Delta(w)$ and set

$$\phi_1 = \sum_{w \in \mathbf{G}(n+r)} \phi(x_{w_1 \dots w_r}) \chi_{\Delta(w)}.$$

Proposition 28 yields the required properties of ϕ_1 (note that we sum over all good words of length $n+r$ in order to be able to apply the Proposition).

We set $\phi_2 = (\phi - \phi_1)\chi_{G(n+r)}$ and $\phi_3 = \phi\chi_{\Delta(B(n+r))}$. The estimate for ϕ_2 is satisfied by the definition of a Hölder function.

Finally, we have

$$\|\phi_3\|_{L_2}^2 = E(|\phi\chi_{\Delta(B(n))}|^2),$$

whence, by Hölder's inequality, using the estimate (28), we obtain the desired estimate for ϕ_3 , and the Proposition is proved completely.

Proposition 29 with $\theta = 1/3$ yields Theorem 4.

We now complete the proof of Theorem 7.

For a word $w \in W_{\mathcal{A},B}$, $|w| = 2n+1$, $w = w_1 \dots w_{2n+1}$, denote $C^{[-n,n]}(w) = \{\omega \in \Omega_{\mathcal{A},B}^{\mathbb{Z}} : \{\omega_{-n} = w_1, \dots, \omega_n = w_{2n+1}\}\}$ and set $\overline{\Delta}(w) = \overline{\Phi}^{-1}C^{[-n,n]}(w)$. Denote by $\mathcal{B}_{[-n,n]}$ the sigma-algebra generated by $\overline{\Delta}(w)$ for all $w \in W_{\mathcal{A},B}$.

Also, for $\epsilon > 0$, denote

$$\overline{\Delta}_\epsilon = \{(\lambda, h, a, \pi) \in \overline{\Delta}(\mathcal{R}) : \lambda \in \Delta_\epsilon\}.$$

Again, we shall prove the Theorem for a slightly larger class of functions.

Say that a function $\phi : \overline{\Delta}(\mathcal{R}) \rightarrow \mathbb{R}$ is *weakly l, α -Hölder* if the following holds. Let k be a positive integer, and let $w \in \mathcal{W}_{\mathcal{A}, B}$, $|w| \leq 2l + 1$ be such that $\overline{\Delta}(w) \subset \Delta_{\exp(-k)}$. Then there exists a constant $C(\phi)$ such that for any $(\lambda, h, a, \pi), (\lambda', h', a', \pi) \in \overline{\Delta}(w)$, we have

$$|\phi(\lambda, h, a, \pi) - \phi(\lambda', h', a', \pi)| \leq Ckd((\lambda, h, a, \pi), (\lambda', h', a', \pi))^\alpha.$$

The smallest such C for a given ϕ will be denoted $C_{l, \alpha}^{weak}(\phi)$. Clearly, if ϕ is Hölder with exponent α , then it is also *weakly l, α -Hölder* for any l and $C_{l, \alpha}^{weak}(\phi) \leq C_\alpha(\phi)$.

Denote by $\overline{G}(2n+1)$ the union of all $\overline{\Delta}(w)$ for good w , by $\overline{B}(2n+1)$ the complement of $\overline{G}(2n+1)$.

Proposition 30 *Let $\theta \in \mathbb{R}$, $0 < \theta < 1$. Let $p > 2$ and $\alpha > 0$. There exist constants C_{101}, C_{102} , such that the following is true for any r and any $n \geq r^2$.*

Let $l \leq r$. Let $\phi \in L_p(\overline{\Delta}(\mathcal{R}) \cdot \mathcal{V})$ be weakly l, α -Hölder and satisfy $\phi \geq 1$. Then there exist functions ϕ_1, ϕ_2, ϕ_3 such that

1. $\phi = \phi_1 + \phi_2 + \phi_3$.
2. ϕ_1 is $\mathcal{B}_{[-n, n]}$ -measurable and supported on $\overline{G}(2n+1)$.
3. $|\phi_2| \leq C_{101}C_\alpha(\phi) \exp(-r^{(1-\theta)/2} + r^\theta)$.
4. $|\phi_3|_{L_2} \leq C_{102} \exp(-r^{(1-\theta)/2}) \|\phi\|_{L_p}$.

For any good w , $|w| = 2n+1$, take an arbitrary point x_w in $\overline{\Delta}(w)$. Set

$$\phi_1 = \sum_{w \in \overline{G}(2n+1)} \phi(x_w) \chi_{\overline{\Delta}(w)},$$

$$\phi_2 = (\phi - \phi_1) \cdot \chi_{\overline{B}(2n+1)},$$

$$\phi_3 = \phi \cdot \chi_{\overline{B}(2n+1)},$$

and the Proposition is proved.

Proposition 30 with $\theta = 1/3$ yields Theorem 7.

It remains to establish the Central Limit Theorem for the flow P^t . Consider the special function $\tilde{\tau}$ of the flow P^t over the transformation $\mathcal{F}$. Note that $\tilde{\tau}(\lambda, h, a, \pi)$ only depends on (λ, π) . Consider the restriction of $\tilde{\tau}$ on a cylinder of the form $\Delta(w_1)$, $w_1 \in \mathcal{A}$. Then there exist distinct $j(1), \dots, j(l) \in \{1, \dots, m\}$ such that

$$\tilde{\tau}(\lambda, \pi) = \log(\lambda_{j(1)} + \lambda_{j(2)} + \dots + \lambda_{j(l)}),$$

which shows that the function $\tilde{\tau}$, restricted to an arbitrary $\Delta(w_1)$ is Lipschitz with respect to the Hilbert metric on $\Delta(\mathcal{R})$.

Now for a Hölder ϕ consider the function

$$\tilde{\phi}(x) = \int_0^{\tilde{\tau}(x)} \phi(P^t x).$$

For any $k > 1$, if $(\lambda, \pi) \in \Delta_{\exp(-k)}$, then, by definition, $\tilde{\tau}(\lambda, \pi) \leq k$. Therefore, if ϕ is Hölder of exponent α , then $\tilde{\phi}$ is weakly $1, \alpha$ -Hölder.

It is easy to see that $\tilde{\tau}(\lambda, \pi) \in L_r(\Delta(\mathcal{R}), \nu)$ for any $r > 1$, whence, if $\phi \in L_p(\Omega_0(\mathcal{R}), \mu_{\mathcal{R}})$ for some $p > 2$, then there exists $p' > 2$ such that the function

$$\tilde{\phi}(x) = \int_0^{\tilde{\tau}(x)} \phi(P^t x)$$

satisfies $\tilde{\phi} \in L_{p'}(\mathcal{Y}^{\pm}, \overline{\nu})$.

Therefore, the Theorem of Melbourne and Török [15] implies Theorem 8, the Central Limit Theorem for the flow P^t .

Acknowledgements.

I am deeply grateful to Yakov G. Sinai, who introduced me to interval exchange transformations, explained to me the method of Markov approximations, and encouraged me in every possible way as the work progressed (more importantly, when it did not).

I am deeply grateful to Alexander Eskin, who suggested to me the problem of the decay of correlations for the induction map.

I am deeply grateful to Giovanni Forni who introduced me to Teichmüller theory.

I am deeply grateful to Corinna Ulcigrai and Pavel Batchourine for their suggestions that have been of invaluable help to me.

I am deeply grateful to Jayadev Athreya, Valdo Durrleman, Charles L. Fefferman, Boris M. Gurevich, Carlangelo Liverani, Michael Ludkovski, Ian Melbourne, Klaus Schmidt, Andrei Török and Anton V. Zorich for useful discussions.

Part of this work was done at The Erwin Schrödinger Institute in Vienna, at The Institute of Mathematics "Guido Castelnuovo" of the University of Rome "La Sapienza", and at the CIRM-IML in Marseille. I am deeply grateful to these institutions for their hospitality.

References

- [1] William Veech, Gauss measures for transformations on the space of interval exchange maps, *Annals of Mathematics*, 15(1982), 201–242.
- [2] W.Veech, Interval exchange transformations. *J. Analyse Math.* 33 (1978), 222–272.

- [3] Veech, William A. Projective Swiss cheeses and uniquely ergodic interval exchange transformations. *Ergodic theory and dynamical systems*, I (College Park, Md., 1979–80), pp. 113–193, *Progr. Math.*, 10, Birkhuser, Boston, Mass., 1981.
- [4] Anton Zorich, Finite Gauss measure on the space of interval exchange transformations. Lyapunov exponents. *Ann. Inst. Fourier (Grenoble)* 46 (1996), no. 2, 325–370.
- [5] Oseledets, V. I. The spectrum of ergodic automorphisms. (Russian) *Dokl. Akad. Nauk SSSR* 168 1966 1009–1011.
- [6] G. Rauzy, Échanges d’intervalles et transformations induites. *Acta Arith.* 34, (1979), no. 4, 315–328.
- [7] M. I. Gordin, The central limit theorem for stationary processes, *Dokl. Akad. Nauk SSSR* 188 (1969), 739–741.
- [8] Carlangelo Liverani, Central limit theorem for deterministic systems. *International Conference on Dynamical Systems (Montevideo, 1995)*, 56–75.
- [9] M. Keane, Interval exchange transformations, *Math. Zeitschrift*, 141(1975), 25 – 31.
- [10] Giovanni Forni, Deviation of ergodic averages for area-preserving flows on surfaces of higher genus. *Ann. of Math. (2)*, 155 (2002), no. 1, 1–103.
- [11] Lai-Sang Young, Recurrence times and rates of mixing. *Israel Journal of Mathematics*, 110(1999), 153–188.
- [12] Véronique Maume-Deschamps, Projective metrics and mixing properties on towers, *Trans. Amer. Math. Soc.*, 353 (2001), no.8, 3371–3389.
- [13] Sinai, Ya. G. Gibbs measures in ergodic theory. *Uspehi Matematicheskikh Nauk* 27 (1972), no. 4(166), 21–64.
- [14] Bunimovich, L. A.; Sinai, Ya. G. Statistical properties of Lorentz gas with periodic configuration of scatterers. *Comm. Math. Phys.* 78 (1980/81), no. 4, 479–497.
- [15] I. Melbourne, A. Torok, Statistical limit theorems for suspension flows, to appear in *Israel Journal of Mathematics*.
- [16] Kolmogorov, A. N. A local limit theorem for classical Markov chains. *Izvestiya Akad. Nauk SSSR. Ser. Mat.* 13, (1949). 281–300.
- [17] Marcelo Viana, *Stochastic Dynamics of Deterministic Systems*.
- [18] Caroline Series, The modular surface and continued fractions. *J. London Math. Soc. (2)* 31 (1985), no.1, 69–80.

- [19] Caroline Series, Geometrical Markov coding of geodesics on surfaces of constant negative curvature. *Ergodic theory and dynamical systems* 6 (1986), no. 4, 601-625.
- [20] Kerckhoff, S. P. Simplicial systems for interval exchange maps and measured foliations. *Ergodic Theory Dynam. Systems* 5 (1985), no. 2, 257–271.
- [21] H. Masur, Interval exchange transformations and measured foliations. *Ann. of Math. (2)* 115 (1982), no. 1, 169–200.
- [22] Doob, J. L. *Stochastic processes*. John Wiley and Sons, New York, 1990.
- [23] V.A. Rokhlin, Exact endomorphisms of a Lebesgue space, *Izv. Akad. nauk SSSR Ser. Mat.* 25 (1961) 499-530.
- [24] V.A. Rokhlin, New progress in the theory of transformations with invariant measure. *Uspehi Mat. Nauk*, 15 no. 4, 3– 26.
- [25] Ya.G. Sinai, *Topics in ergodic theory*, Princeton University Press, 1994.
- [26] M.Kontsevich, A.Zorich, Connected components of the moduli spaces of Abelian differentials with prescribed singularities, *Inventiones mathematicae*, 153(2003), no.3, 631-678.
- [27] I.P. Cornfeld, Ya. G. Sinai, S.V. Fomin, *Ergodic theory*, “Nauka”, Moscow, 1980.
- [28] Hubbard, John; Masur, Howard, Quadratic differentials and foliations. *Acta Math.* 142 (1979), no. 3-4, 221–274.
- [29] M.Kontsevich, A.Zorich, Lyapunov exponents and Hodge theory, “Mathematical Beauty of Physics”, Saclay, 1996.